



**International Conference on
Numerical Analysis and
Approximation Theory**

Fourth Edition

Cluj-Napoca, Romania, September 6–9, 2018

**BOOK OF ABSTRACTS
AND PROGRAM**

This edition is organized jointly with
Tiberiu Popoviciu Institute of Numerical Analysis,
Romanian Academy; www.ictp.acad.ro

Honorary Chairs



Professor GHEORGHE COMAN



Professor PETRU BLAGA

The meeting is devoted to aspects of approximation of functions, integral and differential operators, linear approximation processes, splines, numerical analysis, statistics, stochastic processes, wavelets.

Scientific Committee:

Octavian Agradini (Romania)

Francesco Altomare (Italy)

George Anastassiou (USA)

Carlo Bardaro (Italy)

Emil Căținaș (Romania)

Teodora Căținaș (Romania)

Ioan Gavrea (Romania)

Heiner Gonska (Germany)

Mircea Ivan (Romania)

Giuseppe Mastroianni (Italy)

Gradimir Milovanović (Serbia)

Ioan Rașa (Romania)

Nicolae Suciu (Romania)

Gancho Tachev (Bulgaria)

Webpage: math.ubbcluj.ro/naat2018/

E-mail: naat2018@math.ubbcluj.ro

Invited speakers:

Francesco Altomare (*Italy*)

Carlo Bardaro (*Italy*)

Maria Garrido-Atienza (*Spain*)

Heiner Gonska (*Germany*)

Aaron Melman (*USA*)

Gradimir Milovanović (*Serbia*)

Cihan Orhan (*Turkey*)

Wolfgang L. Wendland (*Germany*)

Organizing Committee:

Octavian Agratini
Teodora Căținaș
Ioana Chiorean
Hannelore Lisei
Sanda Micula
Natalia Roșca
Ágoston Róth
Ildikó Somogyi
Anna Soós (*Vice-rector*)
Radu Trîmbițaș

Volunteers: Anamaria Biriș, Ioana Șomîtcă
Special thanks to Diana Șotropa.

- ◆ The organizers would like to acknowledge the substantial support received from **the Rectorate of Babeș-Bolyai University**, for which they are grateful.
- ◆ This conference was partly funded by grant 39/20.08.2018 of **the National Authority for Scientific Research/the Ministry of Research and Innovation** and sponsored by **the Association of Mathematicians and Computer Scientists of Cluj (AMIC)**.

CONTENTS

Conference schedule	8
List of participants	14
Abstracts of plenary talks	18
Abstracts of contributed talks	32
Addresses of speakers	121

Program of the Registration Desk

Department of Mathematics Office (Room 137), 1st floor,
main building of the Babeş-Bolyai University,
1, M. Kogălniceanu St.

Wednesday, September 5: 17:00 - 20:00

Thursday, September 6: 8:30 - 13:30; 14:30 - 16:30

Friday, September 7: 8:30 - 13:30; 14:30 - 16:30

Saturday, September 8: 8:30 - 13:30

CONFERENCE SCHEDULE

Thursday, September 6

9⁰⁰–9¹⁰ OPENING CEREMONY (Room A)

Plenary Talks

	<i>Chairman: Gradimir Milovanović</i>
9 ¹⁰ -10 ⁰⁰	Francesco Altomare
10 ⁰⁰ -10 ⁵⁰	Wolfgang Wendland

10⁵⁰-11²⁰ COFFEE BREAK

Notes.

- 1) All plenary talks are given in Room A.
- 2) Room A is *N. Iorga* Room and Room B is *T. Popoviciu* Room, both located at the 1st floor of the main building of the Babeş-Bolyai University, 1, M. Kogălniceanu St.
- 3) Please check daily the updated program posted on the doors of the rooms.

Contributed Talks

	Room A	Room B
	<i>Chairman:</i> <i>Heiner Gonska</i>	<i>Chairman:</i> <i>Carlo Bardaro</i>
11 ²⁰ -11 ⁴⁰	Margareta Heilmann	Ralf Rigger
11 ⁴⁰ -12 ⁰⁰	Ioan Raşa	Pawel Wozny
12 ⁰⁰ -12 ²⁰	Ulrich Abel	Antonio Jesus Lopez Moreno
12 ²⁰ -12 ⁴⁰	Sorin Gal	Szilard Csaba Laszlo
12 ⁴⁰ -13 ⁰⁰	Harun Karsli	Adrian Viorel

13⁰⁰ – 15⁰⁰ LUNCH BREAK

Plenary Talk

	<i>Chairman: Francesco Altomare</i>
15 ⁰⁰ -15 ⁵⁰	Heiner Gonska

15⁵⁰-16²⁰ COFFEE BREAK

Contributed Talks

	Room A	Room B
	<i>Chairman: Maria Garrido-Atienza</i>	<i>Chairman: Margareta Heilmann</i>
16 ²⁰ -16 ⁴⁰	Wilfried Grecksch	Vita Leonessa
16 ⁴⁰ -17 ⁰⁰	Bjorn Schmalfuss	Mirella Cappelletti Montano
17 ⁰⁰ -17 ²⁰	Hans-Jörg Starkloff	Bogdan Gavrea
17 ²⁰ -17 ⁴⁰	Ralf Wunderlich	Ana Maria Acu
17 ⁴⁰ -18 ⁰⁰	Nicolae Suciu	Carmen Muraru
18 ⁰⁰ -18 ²⁰	Markus Dietz	Voichita Radu

Friday, September 7

Plenary Talks

	<i>Chairman: Wolfgang Wendland</i>
9 ⁰⁰ -9 ⁵⁰	Gradimir Milovanović
9 ⁵⁰ -10 ⁴⁰	Carlo Bardaro

10⁴⁰-11¹⁰ COFFEE BREAK

Contributed Talks

	Room A	Room B
	<i>Chairman:</i> <i>Harun Karsli</i>	<i>Chairman:</i> <i>Ioan Raşa</i>
11 ¹⁰ -11 ³⁰	Radu Miculescu	Dorian Popa
11 ³⁰ -11 ⁵⁰	Ioan Gavrea	Rodrigo Véjar Asem
11 ⁵⁰ -12 ¹⁰	Radu Păltănea	Michal Veselý
12 ¹⁰ -12 ³⁰	Emre Tas	Emil Cătinaş
12 ³⁰ -12 ⁵⁰	Gumrah Uysal	Radu Trîmbiţaş

12⁵⁰ – 15⁰⁰ LUNCH BREAK

Plenary Talk

	<i>Chairman: Wilfried Grecksch</i>
15 ⁰⁰ -15 ⁵⁰	Cihan Orhan

15⁵⁰-16²⁰ COFFEE BREAK

Contributed Talks

	Room A	Room B
	<i>Chairman:</i> <i>Aaron Melman</i>	<i>Chairman:</i> <i>Ulrich Abel</i>
16 ²⁰ -16 ⁴⁰	Călin Gheorghiu	Heinz-Joachim Rack
16 ⁴⁰ -17 ⁰⁰	Petr Hasil	Robert Vajda
17 ⁰⁰ -17 ²⁰	Nicușor Minculete	Dan Miclaus
17 ²⁰ -17 ⁴⁰	Flavian Georgescu	Mehmet Unver
17 ⁴⁰ -18 ⁰⁰	Daniel Pop	Mohd Ahasan
18 ⁰⁰ -18 ²⁰	Flavius Pătrulescu	Adonia Opreș

19⁰⁰ CONFERENCE DINNER

Grand Hotel Napoca, 1, Octavian Goga St.

web: <http://hotelnapoca.ro/>

Saturday, September 8

Plenary Talks

	<i>Chairman: Cihan Orhan</i>
9 ⁰⁰ -9 ⁵⁰	Maria Garrido-Atienza
9 ⁵⁰ -10 ⁴⁰	Aaron Melman

10⁴⁰-11¹⁰ COFFEE BREAK

Contributed Talks

	Room A	Room B
	<i>Chairman:</i> <i>Ioan Gavrea</i>	<i>Chairman:</i> <i>Mircea Ivan</i>
11 ¹⁰ -11 ³⁰	Alexandru Mitrea	Marius Birou
11 ³⁰ -11 ⁵⁰	Julian Dimitrov	Silviu Urziceanu
11 ⁵⁰ -12 ¹⁰	Maria Crăciun	Ildiko Somogy
12 ¹⁰ -12 ³⁰	Diana Otrocol	Vicuta Neagos
12 ³⁰ -12 ⁵⁰	Alina Baias	Larisa Cheregi
12 ⁵⁰ -13 ¹⁰	Tugba Yurdakadim	Augusta Ratiu

Sunday, September 9

9⁰⁰ EXCURSION

LIST OF PARTICIPANTS

Participants	Page
Ulrich Abel	32
Ana Maria Acu	35
Mohd Ahasan	36
Francesco Altomare	18
Alina Ramona Baias	38, 82
Carlo Bardaro	20
Marius Birou	39
Mirella Cappelletti Montano	40
Emil Căținaș	42
Ganna Chekhanova	46
Larisa Cheregi	43
Maria Crăciun	44
Madalina Dancs	38
Markus Dietz	46
Julian Dimitrov	47
Sorin Gal	49
Maria Garrido-Atienza	22
Bogdan Gavrea	51
Ioan Gavrea	52, 75
Flavian Georgescu	53

Participants	Page
Călin Gheorghiu	54
Heiner Gonska	23
Wilfried Grecksch	55
Petr Hasil	56
Margareta Heilmann	57
Laura Hodis	76
Sever Hodis	38
Harun Karsli	59
Davod Khojasteh Salkuyeh	61
Szilárd Csaba László	63
Vita Leonessa	64
Antonio Jesus Lopez Moreno	66
Alexandra-Ioana Măduța	76
Aaron Melman	24
Dan Miclăuș	68
Radu Miculescu	69
Gradimir V. Milovanović	26
Nicusor Minculete	70
Alexandru Mitrea	71
Camelia Liliana Moldovan	77
Carmen Violeta Muraru	73
Vicuta Neagoș	74, 43
Adonia Augustina Opreș	75

Participants	Page
Cihan Orhan	28
Diana Otrocol	76
Radu Păltănea	77
Flavius Pătrulescu	78
Ion Păvăloiu	42
Daniel Pop	80
Dorian Popa	82
Heinz-Joachim Rack	84
Voichița Adriana Radu	85
Ioan Rașa	87, 57
Augusta Rațiu	89
Ralf Rigger	90
Björn Schmalfuß	92
Andra Silaghi	74
Ildikó Somogyi	93
Hans-Jörg Starkloff	95, 46
Nicolae Suci	96
Emre Taş	97, 120
Radu Trîmbițaș	98
Mehmet Unver	99
Silviu-Aurelian Urziceanu	101
Gumrah Uysal	103
Robert Vajda	106

Participants	Page
Rodrigo Vjar Asem	108
Michal Vesel	110
Adrian Viorel	112
Wolfgang Wendland	30
Pawel Wozny	116
Ralf Wunderlich	118
Tugba Yurdakadim	120, 97

Note: In the sequel the symbol * designates the author giving the talk.

PLENARY TALKS

POSITIVE APPROXIMATION PROCESSES AND INITIAL-BOUNDARY VALUE DIFFERENTIAL PROBLEMS

Francesco Altomare

Department of Mathematics, University of Bari, Italy
[francesco.altomare@uniba.it]

MSC 2010: 41A36, 47D06, 47D07, 35K65, 35B65

Keywords: Approximation by positive operators, positive C_0 -semigroup of operators, initial boundary-value differential problem.

The talk will be centered about a topic concerning three interrelated subjects: positive approximating operators, positive C_0 -semigroups of operators and initial-boundary value evolution problems.

The main aim is to discuss a series of results concerning those sequences $(L_n)_{n \geq 1}$ of bounded linear operators on a Banach space E whose iterates converge to a C_0 -semigroup $(T(t))_{t \geq 0}$ of operators on E .

To such a semigroup it is naturally associated its infinitesimal generator $A : D(A) \rightarrow E$ which, in turn, gives rise to an abstract Cauchy problem (initial-boundary value problem) whose solutions can be given, at least from a theoretical point of view, by the semigroup itself.

Thus, if it is possible to determine the operator A and its domain $D(A)$, then the initial sequence $(L_n)_{n \geq 1}$ becomes the key tool to approximate and to study (especially, from a qualitative point of view) the solutions of the Cauchy problem.

The principal ideas and some of the more recent results on such functional analytic approach to study these kinds of problems, will be discussed in the context of continuous function spaces by also assuming that the operators $L_n, n \geq 1$, are positive. Moreover, particular attention will be devoted to the important case when the approximating operators are constructively generated by a given positive linear operator $T : C(K) \rightarrow C(K)$ which, in turn, allows to determine the differential operator $(A, D(A))$ as well, K being a compact subset of \mathbb{R}^d , $d \geq 1$, having non-empty interior.

Initial-boundary value evolution problems corresponding to these particular settings, occur, for instance, in the study of diffusion problems arising from different areas such as biology, mathematical finance and physics.

For more details and for several other aspects related to the above outlined theory, the reader is referred to the monograph [1].

REFERENCES

- [1] F. Altomare, M. Cappelletti Montano, V. Leonessa and I. Raşa, *Differential Operators, Markov Semigroups and Positive Approximation Processes Associated with Markov Operators*, de Gruyter Series Studies in Mathematics, Vol. 61, De Gruyter, Berlin, 2014.

PALEY-WIENER THEOREMS FOR MELLIN TRANSFORMS AND THE EXPONENTIAL SAMPLING

Carlo Bardaro

*Department of Mathematics and Computer Sciences,
University of Perugia, Italy*
[carlo.bardaro@unipg.it]

MSC 2010: 26D10, 30D20, 44A05

Keywords: Mellin transforms, Paley-Wiener spaces, Bernstein spaces, band-limited functions, polar-analytic functions.

The exponential sampling formula, introduced by a group of physicists and engineers during the end of seventieth, was studied in a rigorous form, through the Mellin transform theory, by P.L. Butzer and S. Jansche during the nineties. This formula is formally equivalent to the classical Shannon sampling formula of signal analysis, valid for Fourier bandlimited functions. Indeed, by a formal change of variable and change of function it is possible to obtain one formula from the other. However, this equivalence is only formal. Indeed, the structure of the Paley-Wiener space of all continuous functions in $L^2(\mathbb{R})$ which are bandlimited, is characterized by the famous Paley-Wiener theorem of Fourier analysis which states that a (Fourier) bandlimited function has an extension to the complex field as an entire function of exponential type (the Bernstein space). In Mellin transform setting this is not true. Indeed, it is shown that a (non trivial) Mellin bandlimited function cannot be extended as an entire function over \mathbb{C} . As proved in [1], it has an extension as an analytic function to the Riemann

surface of the (complex) logarithm with certain exponential type conditions (Mellin-Bernstein spaces). In this talk, we discuss some further version of the Paley-Wiener theorem in Mellin setting, which avoid the use of Riemann surfaces and analytic branches, employing a new concept of analyticity (see [2]). Another version characterizes the space of all the functions whose Mellin transform decays exponentially at infinity involving Hardy-type spaces (see [3]). Applications to exponential sampling theory are described.

REFERENCES

- [1] C. Bardaro, P.L. Butzer, I. Mantellini, G. Schmeisser, *On the Paley-Wiener theorem in Mellin transform setting*, J. Approx. Theory, 207 (2016), pp. 60-75.
- [2] C. Bardaro, P.L. Butzer, I. Mantellini, G. Schmeisser, *A fresh approach to the Paley-Wiener theorem for Mellin transforms and the Mellin-Hardy spaces*, Math. Nachr., 290 (2017), pp. 2759-2774.
- [3] C. Bardaro, P.L. Butzer, I. Mantellini, G. Schmeisser, *A generalization of the Paley-Wiener theorem for Mellin transforms and metric characterization of function spaces*, Fractional Calculus and Applied Analysis, 20 (2017), pp. 1216-1238.

STOCHASTIC DIFFERENTIAL EQUATIONS DRIVEN FRACTIONAL BROWNIAN MOTION

María J. Garrido-Atienza

*Departamento de Ecuaciones Diferenciales y Análisis
Numérico, Universidad de Sevilla, Spain*

[mgarrido@us.es]

In this talk we are concerned with the study of the existence and uniqueness of solutions of stochastic (partial) differential equations driven by a fractional Brownian motion (fBm), as well as their longtime behavior. We will analyze different approaches and consider both the cases of an fBm with Hurst parameter $H \in (1/2, 1)$ and $H \in (1/3, 1/2]$.

This talk is based on some joint works with Duc Hoang Luu (Max-Planck Institut of Leipzig), A. Neuenkirch (University of Mannheim) and B. Schmalfuss (University of Jena).

CHLODOVSKY-BERNSTEIN POLYNOMIAL OPERATORS - PAST AND PRESENT

Heiner Gonska

University of Duisburg - Essen, Germany

[heiner.gonska@uni-due.de]

The talk will be on a long-neglected polynomial approximation process for continuous functions defined on the real half-line. It was introduced by the Russian mathematician Igor Nikolaevich Chlodovsky. Although the first publication on it appeared in 1937 already, progress on investigating it was interrupted for more than half a century. As Butzer and Karsli said in 2009, this approximation process is not so easy to handle.

Some progress was made over the last ten years. We will present some of these latest developments including very recent results.

EIGENVALUE LOCALIZATION FOR MATRIX POLYNOMIALS

Aaron Melman

*Department of Applied Mathematics, Santa Clara University,
CA, USA*

[amelman@scu.edu]

MSC 2010: 12D10, 15A18, 15A42, 15A54, 26C10, 30C10, 30C15, 47A56, 65F15

Keywords: Bound, localization, eigenvalue, scalar, polynomial, matrix polynomial.

We survey a number of well-known and less well-known results ([5], [6]) on the location of scalar polynomial zeros and their generalization to localization results for the eigenvalues of matrix polynomials. Such polynomials occur in polynomial eigenvalues problems, which can be found in a wide range of engineering applications. We include results for matrix polynomials expressed in generalized bases, covering all classical orthogonal bases, such as Hermite, Legendre, Chebyshev, etc.

Finally, we show how some of the aforementioned results for scalar polynomials lead to extensions of the Eneström-Kakeya theorem for polynomials with positive coefficients.

REFERENCES

- [1] A. Melman, *Bounds for eigenvalues of matrix polynomials with applications to scalar polynomials*, Linear Algebra Appl. 504 (2016), pp. 190-203.

- [2] A. Melman, *Improvement of Pellet's theorem for scalar and matrix polynomials*, C. R. Math. Acad. Sci. Paris, 354 (2016), pp. 859-863.
- [3] A. Melman, *Improved Cauchy radius for scalar and matrix polynomials*, Proc. Amer. Math. Soc., 146 (2018), pp. 613624.
- [4] A. Melman, *Eigenvalue bounds for matrix polynomials in generalized bases*, Math. Comp., 87 (2018), pp. 1935-1948.
- [5] G.V. Milovanović, D.S. Mitrinović, and Th. Rassias, *Topics in polynomials: extremal problems, inequalities, zeros*, World Scientific Publishing Co., Inc., River Edge, NJ, 1994.
- [6] Q.I. Rahman, and G. Schmeisser, *Analytic Theory of Polynomials*, London Mathematical Society Monographs. New Series, 26. The Clarendon Press, Oxford University Press, Oxford, 2002.

SOME CLASSES OF ORTHOGONAL POLYNOMIALS IN THE COMPLEX PLANE AND APPLICATIONS

Gradimir V. Milovanović

Mathematical Institute of the Serbian Academy of Sciences and Arts, Serbia

[gvm@mi.sanu.ac.rs]

MSC 2010: 33C45, 33C47, 65D25, 65D30, 78A30

Keywords: Complex orthogonal polynomials, recurrence relations, zeros, numerical differentiation, quadrature formula.

We consider a few classes of polynomials orthogonal in the complex plane with respect to the Hermitian and Non-Hermitian inner products, as well as some applications of such polynomials.

The first class of such complex polynomials was introduced and studied in [1], [2] and [3]. The inner product is not Hermitian and defined by

$$(f, g) = \int_{\Gamma} f(z)g(z)w(z)(iz)^{-1}dz,$$

where $z \mapsto w(z)$ is a *complex* weight function holomorphic in the half disk $D_+ = \{z \in \mathbb{C} \mid |z| < 1, \operatorname{im} z > 0\}$. Polynomials orthogonal on the radial rays in the complex plane is the second class of polynomials in our investigation. This class was introduced in [4] (see also [7], [5], [6]).

Beside some analysis of such kinds of orthogonality and an electrostatic interpretation of zeros of the polynomials on the radial rays, we give several applications of these

polynomials in numerical integration and numerical differentiation.

REFERENCES

- [1] W. Gautschi, G.V. Milovanović, *Polynomials orthogonal on the semicircle*, J. Approx. Theory, 46 (1986), pp. 230–250.
- [2] W. Gautschi, H.J. Landau, G.V. Milovanović, *Polynomials orthogonal on the semicircle. II*, Constr. Approx., 3 (1987), pp. 389–404.
- [3] G.V. Milovanović, *Complex orthogonality on the semicircle with respect to Gegenbauer weight: theory and applications*, In: Topics in Mathematical Analysis (Th. M. Rassias, ed.), pp. 695–722, Ser. Pure Math., 11, World Sci. Publ., Teaneck, NJ, 1989.
- [4] G.V. Milovanović, *A class of orthogonal polynomials on the radial rays in the complex plane*, J. Math. Anal. Appl., 206(1997), pp. 121–139.
- [5] G.V. Milovanović, *Orthogonal polynomials on the radial rays and an electrostatic interpretation of zeros*, Publ. Inst. Math. (Beograd) (N.S.), 64(78)(1998), 53–68.
- [6] G.V. Milovanović, *Orthogonal polynomials on the radial rays in the complex plane and applications*, Rend. Circ. Mat. Palermo (2) Suppl., 68(2002), 65–94.
- [7] G.V. Milovanović, P.M. Rajković, Z.M. Marjanović, *Zero distribution of polynomials orthogonal on the radial rays in the complex plane*, Facta Univ. Ser. Math. Inform., 12(1997), 127–142.

THE SCOTTISH CAFE AND STATISTICAL CONVERGENCE

Cihan Orhan

University of Ankara, Turkey

[Cihan.Orhan@science.ankara.edu.tr]

Many valuable problems and solutions are originated in a cafe so-called "Scottish Cafe" in Lviv around 1930's. These problems are listed in the well-known book *The Scottish Book, Mathematics from the Scottish Cafe*. The participants of this informal meetings includes well-known mathematicians such as Stefan Banach, Hugo Steinhaus, Stanislaw Mazur, W. Orlicz, J.P. Schauder, M. Kac, S. Kacmarz, S. Saks, S. Ulam etc.

Here is the one problem posed by S. Mazur (July 22, 1935):

"A sequence (x_n) is asymptotically convergent to L if there exists a subsequence of density one convergent to L . In the domain of all sequences this notion of convergence is not equivalent to any Toeplitz (regular method). How is it in the domain of bounded sequences?" Mazur also made some comments that indicates the solution of this problem is negative (see, *The Scottish Book*, Second Edition, page 55, problem 5).

In our talk we will provide a positive answer to Mazur's problem and his claim has to be false. In order to prove our result we first note that the notion of asymptotic convergence of Mazur is equivalent to the notion of statistical convergence. We show that statistical convergence is always boundedly (as well as over the space of uniformly

integrable sequences) equivalent to a nonnegative regular matrix method which is boundedly multiplicative. We also characterize the set of bounded multipliers of multiplicative methods. This talk is mainly based on my joint paper with M.K. Khan (Matrix characterization of A -statistical convergence; J. Math. Anal. Appl. 335 (2007), 406-417).

ON NEUMANN'S METHOD AND DOUBLE LAYER POTENTIALS

Wolfgang. L. Wendland

IANS & Simtech University Stuttgart, Germany
[wendland@mathematik.uni-stuttgart.de]

MSC 2010: 31A10, 45F15, 65N30

Keywords: Boundary integral equations, boundary element methods.

Neumann's classical integral equations with the double layer boundary potential is considered on different spaces of boundary charges such as continuous data, L^2 and energy trace spaces on the domain's boundary for interior and exterior boundary value problems of elliptic partial differential equations. Corresponding known results for different classes of boundaries are discussed in view of collocation and Galerkin boundary element methods.

REFERENCES

- [1] O. Steinbach, W.L. Wendland, *On C. Neumann's method for second-order elliptic systems in domains with non-smooth boundaries*, J. Math. Anal. Appl., 262 (2001), pp. 733–748.
- [2] M. Costabel, *Some historical remarks on the positivity of boundary integral operators*, In: Boundary Element Analysis (M. Schanz, O. Steinbach eds.), Springer-Verlag Berlin 2007, pp.1–27.
- [3] W.L. Wendland, *On the double layer potential*, Operator Theory: Advances and Appl. 193 (2009), pp. 319–334.

- [4] G.C. Hsiao, O. Steinbach, W.L. Wendland, *Boundary element methods: Foundation and error analysis*, In: Encyclopedia of Computational Mechanics. Second Edition (E. Stein, R. de Borst, T.J.R. Hughes eds.), 2217 John Wiley & Sons Ltd. pp. 1–62.

CONTRIBUTED TALKS

ASYMPTOTIC PROPERTIES AND OPERATOR NORMS OF GAUSS- WEIERSTRASS OPERATORS AND THEIR LEFT QUASI INTERPOLANTS

Ulrich Abel

*Department Mathematik, Naturwissenschaften und
Datenverarbeitung, Technische Hochschule Mittelhessen,
Wilhelm-Leuschner-Strasse 13, 61169 Friedberg, Germany
[ulrich.abel@mnd.thm.de]*

MSC 2010: 41A36, 41A45, 47A30

Keywords: Approximation by positive operators, operator norm.

The Gauß–Weierstraß convolution operators W_n ($n = 1, 2, 3, \dots$) are defined by

$$(W_n f)(x) = \sqrt{\frac{n}{\pi}} \int_{-\infty}^{\infty} f(t) \exp(-n(t-x)^2) dt \quad (1)$$

(see, e.g., [6, SS 5.2.9]). They are positive linear approximation operators which are applicable to the class $L_c(\mathbb{R})$ of all locally integrable real functions f on \mathbb{R} satisfying the growth condition $f(t) = \mathcal{O}(e^{ct^2})$, as $t \rightarrow \pm\infty$, for some $c > 0$, provided that $n > c$. If f is continuous we have

$$\lim_{x \rightarrow \infty} (W_n f)(x) = f(x)$$

uniformly on compact subsets of \mathbb{R} .

The complete asymptotic expansion of $(W_n f)(x)$ as n tends to infinity appears to be a special case of the results [3] on a more general operator defined by Altomare and Milella [5].

In 2014 Sablonnière [7] defined quasi-interpolants of W_n and studied their basic properties including the operator norm for bounded functions with respect to the sup-norm. In particular, he proved asymptotic expansions for polynomials and functions having a bounded higher order derivative on the whole line.

In the first part we report recent results [2] by presenting the complete asymptotic expansions of $(W_n f)(x)$ and the left quasi-interpolant $(W_n^{[r]} f)(x)$ as n tends to infinity, for functions belonging to the class $L_c(\mathbb{R})$ which are assumed to be only locally smooth. The corresponding results for the Favard operators which are the discrete version of W_n can be found in [4, 1].

Finally, in the second part, we consider the operator norms of W_n and $W_n^{[r]}$ when acting on various function spaces.

REFERENCES

- [1] U. Abel, *Asymptotic expansions for Favard operators and their left quasi-interpolants*, Stud. Univ. Babeş-Bolyai Math. 56 (2011), pp. 199–206.
- [2] U. Abel, O. Agratini, R. Păltănea, *A complete asymptotic expansion for the quasi-interpolants of Gauß–Weierstraß operators*, Mediterr. J. Math. (2018) 15:156, <https://doi.org/10.1007/s00009-018-1195-8>
- [3] U. Abel, M. Ivan, *Simultaneous approximation by Altomare operators*, Proceedings of the 6th international conference on functional analysis and approximation theory, Acquafredda di Maratea (Potenza), Italy, Sept. 24–30, 2009,

- Palermo: Circolo Matematico di Palermo, Rend. Circ. Mat. Palermo, Serie II, Suppl. 82 (2010), pp. 177–193.
- [4] U. Abel, P. L. Butzer, *Complete asymptotic expansion for generalized Favard operators*, Constr. Approx. 35 (2012), pp. 73–88.
DOI: 10.1007/s00365-011-9134-y.
- [5] F. Altomare, S. Milella, *Integral-type operators on continuous function spaces on the real line*, J. Approx. Theory 152 (2008), pp. 107–124.
<http://dx.doi.org/10.1016/j.jat.2007.11.002>
- [6] F. Altomare, M. Campiti, *Korovkin-type Approximation Theory and its Applications*, de Gruyter Studies in Mathematics 17, W. de Gruyter, Berlin, New York, 1994.
- [7] P. Sablonnière, *Weierstrass quasi-interpolants*, J. Approx. Theory 180(2014), pp. 32–48.
<http://dx.doi.org/10.1016/j.jat.2013.12.003>

DIFFERENCES OF POSITIVE LINEAR OPERATORS

Ana Maria Acu

*Department of Mathematics and Informatics, Lucian Blaga
University, Sibiu, Romania*

[anamaria.acu@ulbsibiu.ro]

MSC 2010: 41A25, 41A36

Keywords: First modulus of continuity, positive linear operators.

The results obtained are motivated by the recent results which give a solution to a problem proposed by A. Lupuş in [1]. One of the questions raised by him was to give an estimate for

$$B_n \circ \overline{\mathbb{B}}_n - \overline{\mathbb{B}}_n \circ B_n =: U_n - S_n,$$

where B_n are the Bernstein operators and $\overline{\mathbb{B}}_n$ are the Beta operators. We introduce new inequalities for such differences of positive linear operators and their derivatives in terms of moduli of continuity. This is a joint work with Ioan Raşa from Technical University of Cluj-Napoca, Romania.

REFERENCES

- [1] A. Lupuş, *The approximation by means of some linear positive operators*, in Approximation Theory (M.W. Müller et al., eds), Akademie-Verlag, Berlin, 1995, 201-227.

THE DUNKL GENERALIZATION OF STANCU TYPE Q -SZÁSZ-MIRAKJAN-KANTOROVICH OPERATORS AND SOME APPROXIMATION RESULTS

M. Mursaleen and Mohd. Ahasan*

*Department of Mathematics, Aligarh Muslim University,
Aligarh, 202002, India*

[mursaleenm@gmail.com; ahasan.amu@gmail.com]

MSC 2010: 41A25, 41A36, 33C45

Keywords: q -integers, q -exponential functions, q -hypergeometric functions, Dunkl's analogue, Szász operators, Stancu type q -Szász-Mirakjan-Kantorovich operators, rate of convergence, modulus of continuity and Peetre's K -functional.

In this paper, a Dunkl type generalization of Stancu type q -Szász-Mirakjan-Kantorovich positive linear operators of the exponential function is introduced. With the help of well-known Korovkin's theorem, some approximation properties and also the rate of convergence for these operators in terms of the classical and second-order modulus of continuity, Peetre's K -functional and Lipschitz functions are investigated. Further, some approximation results for bivariate Stancu type q -Szász-Mirakjan-Kantorovich operators are obtained.

REFERENCES

- [1] S.N. Bernstein, *Démonstration du théorème de Weierstrass fondée sur le calcul des probabilités*, Commun. Soc. Math. Kharkow, 2(13)(1912), pp. 1-2.

- [2] H.M. Srivastava, M. Mursaleen, A. Alotaibi, Md. Nasiruzzaman and A.A.H. Al-Abied, *Some approximation results involving the q -Szász-Mirakjan-Kantorovich type operators Via Dunkl's generalization*, Math. Meth. Appl. Sci., 40(15)(2017), pp. 5437-5452.
- [3] O. Szász, *Generalization of S . Bernsteins polynomials to the infinite interval*, J. Res. Natl. Bur. Stand., 45(1950), pp. 239-245.
- [4] S. Sucu, *Dunkl analogue of Szász operators*, Appl. Math. Comput., 244(2014), pp. 42-48.

FUNCTIONS WITH CONVEX IMAGES UNDER BERNSTEIN OPERATORS

Alina Baias^{1,*}, Madalina Dancs², Sever Hodis³

^{1,2,3} *Department of Mathematics, Technical University of
Cluj-Napoca, Romania*

[Baias.Alina@math.utcluj.ro], [dancs_madalina@yahoo.com],
[hodissever@gmail.com]

MSC 2010: 26A51, 26D05

Keywords: Bernstein operators, convex function, log-convex function.

The preservation of convexity under the Bernstein operators B_n is exhaustively investigated in literature.

In this talk we present some non-convex functions f for which $B_n f$ is convex. Inequalities for such functions are also discussed.

QUANTITATIVE RESULTS FOR THE ITERATES OF SOME KING TYPE OPERATORS

Marius-Mihai Birou

*Department of Mathematics, Technical University of Cluj
Napoca, Romania*

[Marius.Birou@math.utcluj.ro]

MSC 2010: 41A25, 41A36

Keywords: Iterates, linear positive operators, convergence.

In this article we define the $-q$ variant of some King type operators which fix the functions e_0 and $e_2 + \alpha e_1$, $\alpha > 0$. We study the rates of convergence for the iterates of these operators using the first and the second order modulus of continuity. We show that the convergence is faster in the case of $-q$ operators ($q < 1$) than in the classical case ($q = 1$).

ON THE POSITIVE SEMIGROUPS GENERATED BY FLEMING-VIOT TYPE DIFFERENTIAL OPERATORS

Mirella Cappelletti Montano

Department of Mathematics, University of Bari, Italy
[mirella.cappellettimontano@uniba.it]

MSC 2010: 35K65, 41A36, 47D06, 47D07

Keywords: Bernstein-Durrmeyer operators with Jacobi weights, Fleming-Viot type differential operator, positive semigroup, approximation of semigroups.

Joint work with Francesco Altomare and Vita Leonessa ([1]).

In this talk, we introduce, in the framework of function spaces defined on the d -dimensional hypercube of \mathbb{R}^d , $d \geq 1$, a class of polynomial type positive linear operators, which generalize the Bernstein-Durrmeyer operators with Jacobi weights on $[0, 1]$ ([3, 2]). By means of them, we study some degenerate second-order elliptic differential operators, often referred to as Fleming-Viot type operators, showing that their closures generate positive semigroups both in the space of all continuous functions and in weighted L^p -spaces. In addition, we show that those semigroups are approximated by iterates the above mentioned Bernstein-Durrmeyer operators.

REFERENCES

- [1] F. Altomare, M. Cappelletti Montano and V. Leonessa, *On the positive semigroups generated by Fleming-Viot type differential operators*, Comm. Pure Appl. Anal., to appear.
- [2] H. Berens and Y. Xu, *On Bernstein-Durrmeyer polynomials with Jacobi weights*, in: C. K. Chui (Ed.), *Approximation Theory and Functional Analysis*, Academic Press, Boston, 1991, pp. 25-46.
- [3] R. Păltănea, *Sur un opérateur polynomial défini sur l'ensemble des fonctions intégrables*, Univ. Babeş-Bolyai, Cluj-Napoca, 83-2 (1983), pp. 101-106.

ON A NEWTON-HERMITE-STEFFENSEN TYPE METHOD

Ion Păvăloiu¹, Emil Cătiñaș^{2,*}

^{1,2}*Tiberiu Popoviciu Institute of Numerical Analysis,
Romanian Academy, Cluj-Napoca, Romania*

[pavaloiu@ictp.acad.ro], [ecatinas@ictp.acad.ro]

MSC 2010: 65H05

Keywords: Inverse interpolatory iterative methods, Newton method, local convergence, monotone convergence.

We consider the solving of nonlinear equations in \mathbb{R} , and we introduce an inverse interpolatory iterative method of Hermite type, for which the nodes are given using the Newton method:

$$y_n = x_n - \frac{f(x_n)}{f'(x_n)}$$
$$x_{n+1} = y_n - \frac{f(y_n)}{f'(y_n)} - \frac{[y_n, y_n, x_n; f]f^2(y_n)}{f'(y_n)[y_n, x_n; f]^2}.$$

The convergence order of the resulted method is 5, and the efficiency index is higher than in the case of the Newton and the Steffensen methods.

Under some natural conditions, the generated iterates converge monotonically to the solution, and one may obtain larger convergence sets than the usual attraction balls.

REFERENCES

- [1] I. Păvăloiu, E. Cătiñaș, *On a robust Aitken-Newton method based on the Hermite polynomial*, Appl. Math. Comput., 287-288 (2016), pp. 224-231.

ON THE POMPEIU MEAN-VALUE THEOREM

Larisa Cheregi^{1,*}, Vicuta Neagos²

^{1,2} *Department of Mathematics, Technical University of Cluj-Napoca, Romania*

[larisa.cheregi@math.utcluj.ro],

[vicuta.neagos@math.utcluj.ro]

MSC 2010: 26A24

Keywords: Pompeiu mean-value theorem, compact set, affine support.

We generalize the Pompeiu mean-value theorem by replacing the graph of a continuous function with a compact set.

REFERENCES

- [1] T. Boggio, *Sur une proposition de M. Pompeiu*, *Mathematica*, Timișoara **23**(1948), pp. 101–102.
- [2] M. Ivan, *A note on a Pompeiu-type theorem*, *Mathematical analysis and approximation theory*, Burg, Sibiu (2002), pp. 129–134.
- [3] D. Pompeiu, *Sur une proposition analogue au théorème des accroissements finis*, *Mathematica* **22**(1946), pp. 143–146.
- [4] O.T. Pop, D. Bărbosu, *A mean-value theorem and some applications*, *Didactica Matematica* **31**(1)(2013), pp. 47–50.
- [5] P.K. Sahoo, T. Riedel, *Mean value theorems and functional equations*, World Scientific Publishing Co., Inc., River Edge, NJ (1998).

A BOSE-EINSTEIN CONDENSATE DARK MATTER MODEL TESTED WITH THE SPARC GALACTIC ROTATION CURVES DATA

Maria Crăciun^{1,*} and Tiberiu Harko²

¹ *Tiberiu Popoviciu Institute of Numerical Analysis,
Romanian Academy, Cluj-Napoca, Romania*

² *Department of Physics, Babes-Bolyai University,
Cluj-Napoca, Romania;*

*School of Physics, Sun Yat-Sen University, Guangzhou
510275, People's Republic of China;*

*Department of Mathematics, University College London,
Gower Street, London, WC1E 6BT, United Kingdom*

[craciun@ictp.acad.ro]

MSC 2010: 83F05, 85A40

Keywords: Cosmology, galactic rotation curves, dark matter models.

In the present study we investigate the properties of the galactic rotation curves in the Bose-Einstein Condensate dark matter model, with quadratic self-interaction, by using more than one hundred galaxies from the recently published Spitzer Photometry & Accurate Rotation Curves (SPARC) data.

REFERENCES

- [1] F. Lelli, S.S. McGaugh, J.M. Schombert, *SPARC: Mass models for 175 disk galaxies with Spitzer Photometry and Accurate Rotation Curves*, *The Astronomical Journal*, 152 (2016), article id. 157.

- [2] X. Zhang, M.H. Chan, T. Harko, S.D. Liang, C.S. Leung, *Slowly rotating Bose Einstein condensate galactic dark matter halos, and their rotation curves*, The European Physical Journal C, 78 (2018), pp. 1-20.

ON A STOCHASTIC ARC FURNACE MODEL

Markus Dietz*, Anna Chekhanova,
Hans-Jörg Starkloff

*Faculty of Mathematics and Computer Science, Technische
Universität Bergakademie Freiberg, Saxony, Germany*
[Markus.Dietz@math.tu-freiberg.de]

MSC 2010: 34F05

Keywords: Electric arc furnace, random differential equation, Ornstein-Uhlenbeck process.

One approach of modeling of an electric arc furnace (EAF) is by the power balance equation which results in a nonlinear ordinary differential equation.

In real-world data can be observed that arc-current and arc-voltage vary randomly in time (cf. [1]), for example they oscillate with a randomly time-varying amplitude and a slight shiver. Therefore it is better to model them as stochastic processes and then solve a random differential equation.

Here we want to propose one example for a modulation by using the Ornstein-Uhlenbeck process and present some results which we gained by studying this model.

This is a joint work with Anna Chekhanova and Hans-Jörg Starkloff.

REFERENCES

- [1] D. Grabowski, J. Walczak, M. Klimas *Electric arc furnace power quality analysis on a stochastic arc model*, 978-1-5386-5186-5/18/\$31.00 2018 IEEE.

AN APPROACH TO DEALING WITH UNCERTAINTY IN NUMERICAL MODELS

Julian Dimitrov

Department of Mathematics, University of Mining and Geology, Sofia, Bulgaria
[juldin@abv.bg]

Keywords: Valuation of dependences, stability of numerical algorithms, principle of uncertainty, relative metric.

The article discusses a method for valuation of the functional dependencies applied for estimation of some numerical models. The method for valuation of dependencies is applied similar to "functional stability analysis" method, which is intended to detect instability in numerical algorithms [1].

The uncertainty principle is a fundamental concept [2], [3]. In this paper, we propose a new approach for dealing with such uncertainty, which combines the uncertainties in the values of parameters and transformation uncertainty that describes local degree of correlation at a given point of domain.

For optimal evaluation of the dependencies, we use relative distance in space of input parameters. The used mathematical theory is based on the theory of functional analysis on metric spaces, where the metric gives an estimation of the error [4]. The p -relative distance was introduced by Ren-Cang [5]. In [6] was introduced the main results of M -relative distances.

We use calculations with semi-logarithmic derivative ensuring accordance with the relative distance and we establish relevant properties. A valuation is applied for an analytical expression.

REFERENCES

- [1] N.J. Higham, *Accuracy and Stability of Numerical Algorithms*, SIAM, Philadelphia (2002).
- [2] H.-W. Bandemer, *Mathematics of Uncertainty - Ideas, Methods, Application Problems*, Springer (2006).
- [3] E. Hofer, *The Uncertainty Analysis of Model Results*, Springer (2018).
- [4] A. Mennucci, *On asymmetric distance*, Anal. and Geom. in Metric Spaces (2013)
- [5] R.-C. Li, *Relative perturbation theory I: Eigenvalue and Singular Value Variations*, SIAM J. Matrix Anal. Appl. 19(4) (1998)
- [6] P. Hasto, *A new weighted metric: the relative metric*, I, J. Math. Anal. Appl. (2002).

APPROXIMATION BY MAX-PRODUCT OPERATORS OF KANTOROVICH TYPE

Lucian Coroianu¹ and Sorin G. Gal^{2,*}

^{1,2} *Department of Mathematics and Computer Science,
University of Oradea, Romania*

[lcoroianu@uoradea.ro], [galso@uoradea.ro]

MSC 2010: 41A35, 41A25, 41A20

Keywords: Max-product operators, max-product operators of Kantorovich kind, uniform approximation, shape preserving properties, localization results, max-product Kantorovich-Choquet operators.

We associate to various linear Kantorovich type approximation operators, nonlinear max-product operators for which we obtain quantitative approximation results in the uniform norm, shape preserving properties and localization results.

REFERENCES

- [1] B. Bede, L. Coroianu and S.G. Gal, *Approximation and shape preserving properties of the Bernstein operator of max-product kind*, Intern. J. Math. Math. Sci. vol. **2009**, Article ID 590589, 2009.
- [2] B. Bede, L. Coroianu and S.G. Gal, *Approximation by Max-Product Type Operators*, Springer, 2016. xv+458 pp., ISBN: 978-3-319-34188-0;
- [3] L. Coroianu and S.G. Gal, *Approximation by truncated max-product operators of Kantorovich-type based on generalized (φ, ψ) -kernels*, submitted.

- [4] L. Coroianu and S.G. Gal, *Approximation by max-product operators of Kantorovich-Choquet type based on generalized (φ, ψ) -kernels*, in preparation.
- [5] L. Coroianu and S.G. Gal, *Classes of functions with improved estimates in approximation by the max-product Bernstein perator*, Anal. Appl. **9**(2011), 249-274.
- [6] L. Coroianu and S.G. Gal, *Localization results for the Bernstein max-product operator*, Appl. Math. Comp., **231**(2014) 73-78.
- [7] S.G. Gal, *Shape-preserving approximation by real and complex polynomials*, Birkhäuser, Boston, Basel, Berlin, 2008.

ON A CONVEXITY PROBLEM

Bogdan Gavrea

*Department of Mathematics, Technical University of
Cluj-Napoca, Romania*

[bogdan.gavrea@math.utcluj.ro]

MSC 2010: 26D15, 26D10, 46N30

Keywords: Linear positive operators, convex functions, Bernstein operators.

The work presented here is a continuation of what was done in [2] and it is strongly connected to the work done in [1]. Applications are given for Mirakyan-Favard-Szász, Baskakov and Szász-Schurer type operators.

REFERENCES

- [1] U. Abel, I. Raşa, *A sharpening of a problem on Bernstein polynomials and convex functions*, Math. Inequal. Appl., 21 (2018), pp. 773-777.
- [2] B. Gavrea, *On a convexity problem in connection with some linear operators*, J. Math. Anal. Appl., 461 (2018), pp. 319-332.

ON THE DECOMPOSITION OF SOME LINEAR POSITIVE OPERATORS

Ioan Gavrea

*Department of Mathematics, Technical University,
Cluj-Napoca, Romania*

[ioan.gavrea@math.utcluj.ro]

MSC 2010: 41A35, 41A10, 41A25

Keywords: Linear operators, Beta operators, positive operators.

In this talk we consider the decomposition of some discretely linear positive operators. Our results generalize the results obtained in [1].

REFERENCES

- [1] M. Heilmann, I. Rasa, *On the decomposition of Bernstein operators*, Numer. Funct. Anal. Optimiz., 36 (2015), pp. 72-85.

INVARIANT MEASURES ASSOCIATED TO φ - MAX - IFSs WITH PROBABILITIES

Flavian Georgescu

*Department of Mathematics and Computer Science,
University of Pitești, Romania, Pitești, Argeș, Romania*
[faviu@yahoo.com]

MSC 2010: 28A80, 37C70, 54H20

Keywords: φ -max-contraction, comparison function, iterated function system with probabilities, Markov operator, fixed point, invariant measure.

We prove that the Markov operator associated to an iterated function system consisting of φ -max-contractions with probabilities has a unique invariant measure whose support is the attractor of the system.

REFERENCES

- [1] F. Georgescu, R. Miculescu, A. Mihail, *A study of the attractor of a φ -max-IFS via a relatively new method*, J. Fixed Point Theory Appl., (2018) 20:24, <https://doi.org/10.1007/s11784-018-0497-6>.
- [2] J. E. Hutchinson, *Fractals and self similarity*, Indiana Univ. Math. J., 30 (1981), pp. 713–747.

SPECTRAL COLLOCATION SOLUTIONS TO A CLASS OF NONLINEAR BVPs ON THE HALF LINE

Călin-Ioan Gheorghiu

Tiberiu Popoviciu Institute of Numerical Analysis, Romanian Academy, Cluj-Napoca, Romania

[ghcalin@ictp.acad.ro]

MSC 2010: 65L10, 65L60, 65L70, 65N35

Keywords: Collocation, Laguerre-Gauss-Radau, Kidder problem, nonlinear Schrödinger, radially symmetric solutions.

The existence, uniqueness and regularity of solutions to the boundary value problems

$$\begin{aligned} \frac{1}{p(x)} (p(x) u'(x))' &= q(x) f(x, u(x), p(x) u'(x)), \quad x \in (0, \infty) \\ \alpha u(0) - \beta u'(0) &= r, \quad \lim_{x \rightarrow \infty} u(x) = 0, \end{aligned} \tag{1}$$

are established. We assume $\alpha > 0$, $\beta \geq 0$ and r is a given constant and f , p and $\frac{1}{q}$ are continuous. Our aim is to accurately approximate the solutions of these problems, as well as to some PDEs reducible to (1), by a high order Laguerre-Gauss-Radau collocation method (see [1], Ch. 2).

REFERENCES

- [1] C.I. Gheorghiu, *Spectral Collocation Solutions to Problems on Unbounded Domains*, Casa Cărții de Știință, Cluj-Napoca, 2018 (see also <https://ictp.acad.ro/gheorghiu>)

PARAMETER ESTIMATIONS FOR A LINEAR PARABOLIC FRACTIONAL SPDE WITH JUMPS

Wilfried Grecksch^{1,*}, Hannelore Lisei,
Jens Lueddeckens

¹*Institute of Mathematics, Martin-Luther-University of
Halle-Wittenberg, Germany*
[wilfried.grecksch@mathematik.uni-halle.de]

MSC 2010: 60H15, 62F12, 60G22

Keywords: Stochastic partial differential equations, parameter estimations.

A drift parameter estimation problem is studied for a linear parabolic stochastic partial differential equation driven by a multiplicative cylindrical fractional Brownian motion with Hurst index $h \in]1/2, 1[$ and a multiplicative Poisson process with values in a Hilbert space. Equations are introduced for the Galerkin approximations of the mild solution process. A mean square estimation criterion is used for these equations. It is proved that the estimate is unbiased and weakly consistent for the original problem.

REFERENCES

- [1] I. Cialenko, *Parameter estimation for SPDEs with multiplicative fractional noise*, Stochastics and Dynamics, 10 (2010) no. 4, pp. 561-576.
- [2] J. Lueddeckens, *Fraktale stochastische Integralgleichungen im White-Noise-Kalkül*, Dissertation Martin-Luther-Universität Halle-Wittenberg, April 2017.

HALF-LINEAR DIFFERENTIAL OPERATORS IN OSCILLATION THEORY

Petr Hasil

*Department of Mathematics and Statistics, Faculty of Science,
Masaryk University, Czech Republic*
[hasil@mail.muni.cz]

MSC 2010: 34C10, 34C15

Keywords: Riccati technique, half-linear equation, oscillation theory, oscillation constant, conditional oscillation.

This is a joint work with M. Veselý. We consider half-linear differential equations given by operators with one-dimensional p -Laplacian. The main subject of this talk is to present results concerning the conditional oscillation of such equations, i.e., we find a border value which separates oscillatory equations from non-oscillatory ones and we explicitly determine this borderline (depending on the equations coefficients). For some of the presented results, we refer to [1, 2].

REFERENCES

- [1] P. Hasil, M. Veselý, *Oscillation and non-oscillation of half-linear differential equations with coefficients determined by functions having mean values*, Open Math., 16 (2018), pp. 507–521.
- [2] P. Hasil, M. Veselý, *Oscillation and non-oscillation results for solutions of perturbed half-linear equations*, Math. Methods Appl. Sci., 41 (2018), pp. 3246–3269.

REPRESENTATIONS FOR K -TH ORDER KANTOROVICH MODIFICATIONS OF LINKING OPERATORS

Margareta Heilmann^{1,*}, Ioan Raşa

¹*School of Mathematics and Natural Sciences, University of Wuppertal, Gaußstraße 20, D-42119 Wuppertal, Germany*
[heilmann@math.uni-wuppertal.de]

MSC 2010: 41A36, 41A10, 41A28

Keywords: Linking operators, Baskakov-Durrmeyer type operators, Kantorovich modifications of operators.

In our talk we investigate linking operators acting on the unbounded interval $[0, \infty)$. Linking operators for the Szász-Mirakjan case were defined by Păltănea in [3] and for Baskakov type operators by Heilmann and Raşa in [2]. If the linking parameter ρ is a natural number we present a representation for Kantorovich variants of arbitrary order in terms of the Baskakov and Szász-Mirakjan basis functions, respectively. This leads to a simple proof of convexity properties which also solves an open problem mentioned in [1]. At the end of our talk we present a conjecture concerning a limit to B-splines which is connected to the above mentioned representation.

REFERENCES

- [1] K. Baumann, M. Heilmann, I. Raşa, *Further results for k th order Kantorovich modification of linking Baskakov type operators*, Results Math. 69 (3), (2016), 297-315.
- [2] M. Heilmann, I. Raşa, *k -th order Kantorovich modification of linking Baskakov type operators*, in: Recent Trends in Mathematical Analysis and its Applications, Proceedings of the Conference ICRTMAA 2014, Rorkee, India, December 2014, (ed. P. N. Agrawal et al.), Proceedings in Mathematics & Statistics 143, 229-242.
- [3] R. Păltănea, *Modified Szász-Mirakjan operators of integral form*, Carpathian J. Math., 24(3) (2008), 378-385.

SOME APPROXIMATION PROPERTIES OF URYSOHN TYPE OPERATORS

Harun Karsli

Department of Mathematics, University of Bolu Abant Izzet Baysal, Turkey

[karsli_h@ibu.edu.tr]

MSC 2010: 41A25, 41A35, 47G10, 47H30

Keywords: Urysohn operators, linear positive operators, approximation.

In the present work, our aim is generalization and extension of the theory of interpolation of functions to functionals by means of Urysohn type operators. In accordance with this purpose, we introduce and study a new type of Urysohn type operators. In particular, we investigate the convergence problem for operators that approximate the Urysohn type operator. We construct our operators by using a nonlinear forms of the kernels together with the Urysohn type operator values instead of the sampling values of the function.

REFERENCES

- [1] C. Bardaro, I. Mantellini, *On the reconstruction of functions by means of nonlinear discrete operators*. J. Concr. Appl. Math. 1 (2003), no. 4, pp. 273–285.
- [2] C. Bardaro, J. Musielak, G. Vinti, *Nonlinear Integral Operators and Applications*, De Gruyter Series in Nonlinear Analysis and Applications, Vol. 9, xii + 201 pp., 2003.

- [3] C. Bardaro, G. Vinti, *Urysohn integral operators with homogeneous kernel: approximation properties in modular spaces*, Comment. Math. (Prace Mat.) 42 (2002), no. 2, pp. 145-182.
- [4] C. Bardaro, H. Karsli and G. Vinti, *Nonlinear integral operators with homogeneous kernels: pointwise approximation theorems*, Applicable Analysis, Vol. 90, Nos. 3-4, March-April (2011), pp. 463-474.
- [5] C. Bardaro, H. Karsli and G. Vinti, *On pointwise convergence of linear integral operators with homogeneous kernels*, Integral Transforms and Special Functions, 19(6), (2008), pp. 429-439.
- [6] D. Costarelli, G. Vinti, *Degree of approximation for nonlinear multivariate sampling Kantorovich operators on some functions spaces*, Numer. Funct. Anal. Optim. 36 (2015), no. 8, pp. 964-990.
- [7] D. Costarelli, G. Vinti, *Approximation by nonlinear multivariate sampling Kantorovich type operators and applications to image processing*, Numer. Funct. Anal. Optim. 34 (2013), no. 8, pp. 819-844.
- [8] I.I. Demkiv, *On Approximation of the Urysohn operator by Bernstein type operator polynomials*, Visn. L'viv. Univ., Ser. Prykl. Mat. Inform., Issue 2, pp. 26 - 30.
- [9] H. Karsli, *Approximation by Urysohn type Meyer-König and Zeller operators to Urysohn integral operators*, Results Math. 72 (2017), no. 3, pp. 1571-1583.
- [10] H. Karsli, *Some convergence results for nonlinear singular integral operators*, Demonstratio. Math., Vol. XLVI No 4, (2013), pp. 729-740.
- [11] V.L. Makarov, and I.I. Demkiv, *Approximation of the Urysohn operator by operator polynomials of Stancu type*, Ukrainian Math Journal, 64(3)(2012), pp. 356 - 386.

A NEW METHOD FOR SOLVING COMPLEX SYMMETRIC SYSTEMS OF LINEAR EQUATIONS

Davod Khojasteh Salkuyeh

Department of Mathematics, University of Guilan, Rasht, Iran
[e-mail khojasteh@guilan.ac.ir]

MSC 2010: 65F10

Keywords: Complex, symmetric, convergence, symmetric positive definite.

We consider the system of linear equations $(W + iT)u = b$, where $W, T \in \mathbb{R}^{n \times n}$ are symmetric positive semidefinite matrices with at least one of them being positive definite and $i = \sqrt{-1}$. Letting $b = p + iq$ and $u = x + iy$, we propose a new iterative method for two-by-two block real equivalent form

$$\begin{bmatrix} W & -T \\ T & W \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} p \\ q \end{bmatrix},$$

of the system. Convergence of the proposed method is studied and the numerical results of the method are compared with those of the MHSS [1], the PMHSS [2], the TTSCSP [3], the CRI [4] methods.

REFERENCES

- [1] Z.-Z. Bai, M. Benzi, F. Chen, *Modified HSS iterative methods for a class of complex symmetric linear systems*, Computing 87 (2010), pp. 93-111.

- [2] Z.-Z. Bai, M. Benzi, F. Chen, *On preconditioned MHSS iteration methods for complex symmetric linear systems*, Numer. Algor. 56 (2010), pp. 93-317.
- [3] D.K. Salkuyeh, T.S. Siahkolaei, *Two-parameter TSCSP method for solving complex symmetric system of linear equations*, Calcolo (2018), pp. 55: 8. <https://doi.org/10.1007/s10092-018-0252-9>.
- [4] T. Wang, Q. Zheng, L. Lu, *A new iteration method for a class of complex symmetric linear systems*, J. Comput. Appl. Math., 325 (2017), pp. 188-197.

CONVERGENCE RATES FOR AN INERTIAL ALGORITHM OF GRADIENT TYPE ASSOCIATED TO A SMOOTH NONCONVEX MINIMIZATION

Szilárd Csaba László

*Department of Mathematics, Technical University of
Cluj-Napoca, Romania*

[laszlosziszi@yahoo.com]

MSC 2010: 90C26, 90C30, 65K10

Keywords: Inertial algorithm, nonconvex optimization, Kurdyka-Lojasiewicz inequality, convergence rate.

We investigate an inertial algorithm of gradient type in connection with the minimization of a nonconvex differentiable function. The algorithm is formulated in the spirit of Nesterov's accelerated convex gradient method. We show that the generated sequences converge to a critical point of the objective function, if a regularization of the objective function satisfies the Kurdyka-Lojasiewicz property. Further, we provide convergence rates for the generated sequences and the objective function values formulated in terms of the Lojasiewicz exponent.

ON THE INTERPLAY BETWEEN MARKOV OPERATORS PRESERVING POLYNOMIALS AND CONVEX COMPACT SUBSETS OF \mathbb{R}^d

Vita Leonessa

*Department of Mathematics, Computer Science and
Economics, University of Basilicata, Potenza, Italy*
[vita.leonessa@unibas.it]

MSC 2010: 47B65, 47D07

Keywords: Markov operators, polynomial preserving property, second-order elliptic differential operator, Markov semigroup.

Let K be a convex compact subset of \mathbb{R}^d , $d \geq 1$, with non-empty interior. In this talk we are interesting to all positive linear operators T acting on the space $C(K)$ of continuous functions on K which leave invariant the polynomials of degree at most 1 and which, in addition, map polynomials into polynomials of the same degree m , $m \geq 2$.

In particular, we discuss the existence of operators T when K is a non-trivial strictly convex subset of \mathbb{R}^d , $d \geq 2$, discovering, among other things, a characterization of ellipsoids and balls of \mathbb{R}^d . A discussion of the above polynomial preserving property in the setting of product spaces, as well as for convex convolution products of positive linear operators, is also presented.

Such a polynomial preserving property play a central role in studying the possibility to get a representation/approximation formula for semigroups generated by certain differential operators, associated with T , in terms of

constructively defined linear positive operators associated with the same T (see, e.g. [1, 4]). Anyway, we point out that this is not the only case where a property of this kind is required (see, e.g. [3]). Moreover, in the literature there are many classes of positive linear operators that satisfy it.

All results presented are contained in the joint work [2].

REFERENCES

- [1] F. Altomare, M. Cappelletti Montano, V. L., I. Raşa, *On differential operators associated with Markov operators*, J. Funct. Anal. **266** (2014), pp. 3612–3631.
- [2] F. Altomare, M. Cappelletti Montano, V. L., I. Raşa, *On Markov operators preserving polynomials*, J. Math. Anal. Appl. **415** (2014), pp. 477–495.
- [3] O. Agratini, *A sequence of positive linear operators associated with an approximation process*, Appl. Math. Comput. **269** (2015), pp. 23–28.
- [4] F. Altomare, M. Cappelletti Montano, V. L., I. Raşa, *Elliptic differential operators and positive semigroups associated with generalized Kantorovich operators*, J. Math. Anal. Appl. **458** (2018), pp. 153–173.

DURRMEYER TYPE OPERATORS PRESERVING EXPONENTIAL FUNCTIONS

Vijay Gupta and A.-J. López-Moreno*

*Department of Mathematics, University of Jaén, Campus Las
Lagunillas, 23071-Jaén, Spain*

[ajlopez@ujaen.es]

MSC 2010: 41A25, 41A36

Keywords: Phillips operators, Durrmeyer operators, exponential functions, moments, quantitative results, weighted approximation, asymptotic expansion, Voronovskaja formula.

In this paper (see [5]) we present a modification of certain family of Durrmeyer type operators operators which reproduces two given functions that generalizes many of the Durrmeyer versions that we find in the recent literature [1, 2, 4, 6, 7]. We study the moments and basic approximation properties for these new operators. For this purpose we adapt several already known result to the particular features of the new sequence that we propose here.

REFERENCES

- [1] T. Acar, *Asymptotic formulas for generalized Szász-Mirakyan operators*, Appl. Math. Comput. **263** (2015), 233–239.
- [2] T. Acar, A. Aral, D. Cárdenas-Morales and P. Garrancho, *Szász-Mirakyan type operators which fix exponentials*, Results Math. **72** (2017) no. 3, 1393–1404.

- [3] T. Acar, A. Aral and H. Gonska, *On Szász–Mirakyan operators preserving e^{2ax} , $a > 0$* , Mediterr. J. Math. **14** (2017), no. 1 Art. 6, doi:10.1007/s00009-016-0804-7.
- [4] V. Gupta, *A note on modified Phillips operators*, Southeast Asian Bulletin of Mathematics 34 (2010), 847-851.
- [5] V. Gupta, A.-J. López-Moreno, *Phillips operators preserving arbitrary exponential functions, e^{at} , e^{bt}* , Filomat, to appear.
- [6] V. Gupta and G. Tachev, *On approximation properties of Phillips operators preserving exponential functions*, Mediterr. J. Math. 14 (2017), no. 4, Art. 177.
- [7] G. Tachev, V. Gupta and A. Aral, *Voronovskaja’s theorem for functions with exponential growth*, Georgian Math. J., to appear.

SOME NEW RESULTS CONCERNING THE CLASSICAL BERNSTEIN QUADRATURE FORMULA

Dan Miclăuș

Department of Mathematics and Computer Science, Technical University of Cluj-Napoca, North University Center at Baia Mare, Romania

[dan.miclaus@cunbm.utcluj.ro]

MSC 2010: 41A36, 41A80.

Keywords: Bernstein operator, convex function, divided difference, Popoviciu theorem, remainder term.

In this talk, we intend to highlight an applicative side of the classical Bernstein polynomials, in contrast to the well-known theory of the uniform approximation of functions. We will present some new results concerning the classical Bernstein quadrature formula

$$\int_a^b F(x)dx \approx \frac{b-a}{n+1} \sum_{k=0}^n F\left(a + \frac{k(b-a)}{n}\right),$$

which can be found in [1] and in another recent paper submitted for publication.

REFERENCES

- [1] D. Miclăuș, L. Pișcoran, *A new method for the approximation of integrals using the generalized Bernstein quadrature formula*, Appl. Math. Comput., (accepted for publication 2018).

A NEW ALGORITHM THAT GENERATES THE IMAGE OF THE ATTRACTOR OF A GIFS

Radu Miculescu

*Department of Mathematics and Computer Science,
University of Bucharest, Romania, Academiei Street 14,
010014, Bucharest, Romania*
[miculesc@yahoo.com]

MSC 2010: 28A80, 37C70, 41A65, 65S05, 65P99

Keywords: Generalized infinite iterated function system (GIFS), attractor, deterministic algorithm, grid algorithm.

We provide a new algorithm (called the grid algorithm) designed to generate the image of the attractor of a generalized iterated function system and we compare it with the deterministic algorithm regarding generalized iterated function systems presented by P. Jaros, Ł. Maślanka and F. Strobil in [Algorithms generating images of attractors of generalized iterated function systems, *Numer. Algor.*, 73 (2016), 477-499].

REFERENCES

- [1] P. Jaros, Ł. Maślanka, F. Strobil, *Algorithms generating images of attractors of generalized iterated function systems*, *Numer. Algor.*, 73 (2016), pp. 477-499.
- [2] A. Mihail, R. Miculescu, *Applications of Fixed Point Theorems in the Theory of Generalized IFS*, *Fixed Point Theory Appl.* Volume 2008, Article ID 312876, 11 pages doi: 10.1155/2008/312876.
- [3] A. Mihail, R. Miculescu, *Generalized IFSs on Noncompact Spaces*, *Fixed Point Theory Appl.* Volume 2010, Article ID 584215, 11 pages doi: 10.1155/2010/584215.

ON SEVERAL INEQUALITIES IN AN INNER PRODUCT SPACE

Nicușor Minculete

Transilvania University, Romania
[minculeten@yahoo.com]

MSC 2010: 46C05, 26D15, 26D10

Keywords: Inner product space, Cauchy-Schwarz inequality.

The aim of this presentation is to prove new results related to several inequalities in an inner product space. Among these inequalities we will mention inequality Cauchy-Schwarz's inequality. Also we obtain some applications of these inequalities.

ON THE UNBOUNDED DIVERGENCE OF SOME INTERPOLATORY PRODUCT INTEGRATION RULES

Alexandru Mitrea

Department of Mathematics, Technical University of Cluj-Napoca, Romania

[alexandru.ioan.mitrea@math.utcluj.ro]

MSC 2010: 41A10, 65D32

Keywords: Product integration, Condensation of singularities, superdense set.

Based on some principles of Functional Analysis, this paper emphasizes the topological structure of the set of unbounded divergence for some interpolatory product quadrature formulas on Jacobi and equidistant points of the interval $[-1, 1]$, associated with the Banach space of all s -times continuously differentiable functions and with a weighted Banach space of absolutely integrable functions of order $p > 1$.

REFERENCES

- [1] H. Brass, K. Petras, *Quadrature Theory. The Theory of Numerical Integration on a Compact Interval*, American Mathematical Society, Providence (2011).
- [2] S. Cobzas, I. Muntean, *Condensation of singularities and divergence results in Approximation Theory*, J. Approx. Theory, 31 (1981), pp. 138-153.

- [3] A.I. Mitrea *On the dense unbounded divergence of interpolatory product integration on Jacobi nodes*, *Calcolo*, vol.55, No.1, Art.10 (2018), pp. 1-15.
- [4] I.H. Sloan, W.E. Smith, *Properties of interpolatory product integration rules*, *SIAM J. Numer. Anal.*, 19 (1982), pp. 427-442.

APPROXIMATION PROPERTIES OF CERTAIN BERNSTEIN-STANCU TYPE OPERATORS

Carmen Violeta Muraru

Department of Mathematics and Informatics, Vasile Alecsandri University of Bacau, Romania
[cmuraru@ub.ro]

MSC 2010: 41A36, 41A25

Keywords: Bernstein-Stancu operator, q -integers, rate of convergence, moduli of continuity.

In this paper we introduce and investigate a new operator of Bernstein-Stancu type, based on q polynomials. We study approximation properties for these operators based on Korovkin type approximation theorem and study some direct theorems. Also, the study contains numerical considerations regarding the constructed operators based on Maple algorithms.

This is a joint work with Ana-Maria Acu from Lucian Blaga University of Sibiu, Romania, Ogun Dogru from Gazi University, Turkey and Voichita Adriana Radu from Babes-Bolyai University, Romania.

ON THE FIRST MEAN-VALUE THEOREM FOR INTEGRALS

Vicuta Neagos^{1,*}, Andra-Gabriela Silaghi²

^{1,2}*Department of Mathematics, Technical University of Cluj-Napoca, Romania*

[vicuta.neagos@math.utcluj.ro],

[fun_tastiq@yahoo.com]

MSC 2010: 26A24

Keywords: Positive linear functionals, mean-value theorems, integrals of Riemann and Lebesgue type.

We improve the classical First Mean-value Theorem for Integrals and obtain related results.

REFERENCES

- [1] R.G. Bartle, *A modern theory of integration*, Graduate Studies in Mathematics, vol. 32, American Mathematical Society, Providence, RI (2001).
- [2] P. Khalili, D. Vasiliu, *An extension of the mean value theorem for integrals*, Internat. J. Math. Ed. Sci. Tech. 41(5)(2010), pp. 707–710.
- [3] M. Monea, *An integral mean value theorem concerning two continuous functions and its stability*, Int. J. Anal., Art. ID 894,625, 4 (2015).
- [4] P.K. Sahoo, T. Riedel, *Mean value theorems and functional equations*, World Scientific Publishing Co. Inc., River Edge, NJ (1998).
- [5] S.G. Wayment, *An integral mean value theorem*, Math. Gaz., 54(389) (1970), pp. 300–301.

MODIFIED KANTOROVICH-STANCU OPERATORS (II)

Ioan Gavrea¹, Adonia-Augustina Opriș^{2,*}

^{1,2}*Department of Mathematics, Technical University of Cluj-Napoca, str. Memorandumului, nr. 28, 400144, Cluj-Napoca, Romania*

[ioan.gavrea@math.utcluj.ro], [mate.salaj@yahoo.com]

MSC 2010: 41A25, 41A36

Keywords: Approximation by linear operators, Kantorovich-Stancu operators.

In this paper we introduce a new kind of Bernstein-Kantorovich-Stancu operators. This operators generalizes the operators introduced in the article *Modified Kantorovich operators with better approximation properties* by V. Gupta, G. Tachev, A. M. Acu, published online 18 may 2018 in Springer, Numer. Algor, <https://doi.org/10.1007/s11075-018-0538-7>.

ULAM CONSTANTS FOR FUNCTIONAL EQUATIONS

Laura Hodiş, Alexandra Măduţa,
Diana Otrocol*

*Department of Mathematics, Technical University of
Cluj-Napoca, 28 Memorandumului Street, 400114 Cluj-Napoca,
Romania*

[Diana.Otrocol@math.utcluj.ro]

MSC 2010: 39B82

Keywords: Cauchy, Jensen, Quadratic equations, Hyers-Ulam stability, best constant.

We investigate the best Ulam constants for the Cauchy, Jensen and Quadratic functional equations. Our results are related to those presented in [1, Section 4.2].

REFERENCES

- [1] J. Brzdęk, D. Popa, I. Raşa, B. Xu, *Ulam Stability of Operators*, Mathematical Analysis and its Applications, Series Editor Themistocles M. Rassias, Academic Press, 2018, ISBN: 978-0-12-809829-5, 225 pp.

SCHOENBERG SPLINES WITH KNOTS AT THE ROOTS OF CHEBYSHEV POLYNOMIALS

Radu Păltănea^{1,*} and Camelia Moldovan²

^{1,2} *Department of Mathematics an Informatics, "Transilvania" University of Braşov, Romania*
[radupaltanea@unitbv.ro]

MSC 2010: 41A36

Keywords: Spline functions, Schoenberg operators, degree of approximation.

The paper studies the degree of approximation of continuous functions using the Schoenberg type operators with knots at the roots of Chebyshev polynomials.

REFERENCES

- [1] L. Beutel, H. Gonska, D. Kacso, G. Tachev, *On variation-diminishing Schoenberg operators: new quantitative statements*, Monogr. Academia Ciencias de Zaragoza 20 (2002), pp. 9-58.
- [2] M.J. Marsden, *An identity for spline functions with applications to variation diminishing spline approximation*, J. Approx. Theory 3 (1970), pp. 7-49.
- [3] R. Păltănea, *Approximation Theory Using Positive Linear Operators*, Birhauser, Boston, (2004).
- [4] I.J. Schoenberg, *On variation diminishing approximation methods*, On Numerical Approximation, MRC Symposium (R.E. Langer, ed.), Univ. of Wisconsin Press, Madison, (1959), pp. 249-274.

NATURAL CONVECTION FROM A VERTICAL PLATE EMBEDDED IN A NON-DARCY BIDISPERSE POROUS MEDIUM

Teodor Groșan¹ and Flavius Pătrulescu^{2,*}

¹*Department of Mathematics, Babeș-Bolyai University, Cluj-Napoca, Romania*

²*Tiberiu Popoviciu Institute of Numerical Analysis, Romanian Academy, Cluj-Napoca, Romania*

[tgrosan@math.ubbcluj.ro], [fpatrulescu@ictp.acad.ro]

MSC 2010: 76S99, 76R10, 76M25

Keywords: Bidisperse porous medium, free convection, boundary layer, non-Darcy flow.

A mathematical model for the free convection from a vertical plate placed in a non-Darcy BDPM (bidisperse porous medium) containing inertial terms is proposed. The influence of the inertial parameters on the fluid and heat transfer characteristics is studied. The partial differential equations governing the flow and heat transfer in the f-phase and the p-phase are solved numerically using the `bvp4c` routine from Matlab.

REFERENCES

- [1] D.A. Nield, A.V. Kuznetsov, *Natural convection about a vertical plate embedded in a bidisperse porous medium*, International Journal of Heat and Mass Transfer, 51 (2008), pp. 1658–1664.
- [2] J. Kierzenka, L.F. Shampine, *A BVP solver based on residual control and MATLAB PSE*, ACM Transactions on Mathematical Software, 27 (2001), pp. 299–316.

ON A PARABOLIC PDE

Daniel N. Pop

*Faculty of Engineering, Lucian Blaga University, Sibiu,
Romania*

[nicolae.pop@ulbsibiu.ro,popdaniel31@yahoo.com]

MSC 2010: 35K20, 34B05, 65M12

Keywords: Parabolic equation and systems, boundary value problems, numerical analysis.

Consider the following parabolic PDE

$$\frac{\partial P}{\partial t} = A \frac{\partial P}{\partial x} \left(P \frac{\partial P}{\partial x} \right),$$

with initial condition

$$P(x, 0) = P_0$$

and boundary condition

$$P(0, t) = P_1, \text{ where } \alpha = 1 - \frac{P_0^2}{P_1^2}.$$

The origin of this problem is in the study of a gas flow through a semi-infinite porous medium (see [1]).

We present two solution for this problem: by reduction to a BVP for ODE and by using line method.

REFERENCES

- [1] P.B. Bailey et al, *Nonlinear two-points boundary value problems*, Academic Press, 1968.
- [2] U. Ascher, B. Mateij, R.D. Russel, *Numerical solution of boundary problems for ordinary differential equations*, SIAM, Prentice Hall, 1997.
- [3] C.M. Bender, S.A. Orsag, *Advanced Mathematical Methods for Scientists and Engineers I.-Asymptotic Methods and Perturbation Theory*, New-York, Springer Verlag, 1999.
- [4] D.N. Pop, R.T. Trîmbițaș, *Aproximation method for second order Nonlinear value polylocal problems using B-Spline and Runge Kutta Methods*, N.A.T 2010 Studia univ. Babeș-Bolyai, Mathematica, Vol LVI, Number 2, 2011, pp 515-527.
- [5] L.F. Shampine, I.Gladwell, S. Thomson, *Solving ODE with Matlab*, Cambridge University Press, 2003.

ON THE BEST ULAM CONSTANT OF THE LINEAR DIFFERENTIAL OPERATOR

Alina-Ramona Baias¹, Florina Blaga²,
Dorian Popa^{3,*}

^{1,3}*Department of Mathematics, Technical University of Cluj-Napoca, Romania*

²*Technical University of Cluj-Napoca, Romania*

[Baias.Alina@math.utcluj.ro],

[Blaga.Florina@math.utcluj.ro],

[Popa.Dorian@math.utcluj.ro]

MSC 2010: 34D20, 39B82

Keywords: Ulam stability, Best Ulam constant, linear differential operator.

Let X be a Banach space over \mathbb{C} , $a_1, a_2, \dots, a_n \in \mathbb{C}$ and $D : \mathcal{C}^n(\mathbb{R}, X) \rightarrow \mathcal{C}(\mathbb{R}, X)$ given by the relation

$$Dy = y^{(n)} + a_1 y^{(n-1)} + \dots + a_n y, y \in \mathcal{C}^n(\mathbb{R}, X). \quad (1)$$

Define $\|\cdot\|_\infty : \mathcal{C}^n(\mathbb{R}, X) \rightarrow \overline{\mathbb{R}}$ by

$$\|y\|_\infty = \sup_{x \in \mathbb{R}} \|y(x)\|. \quad (2)$$

The operator D is called Ulam stable if there exists $K \geq 0$ such that for every $\varepsilon > 0$ and every $y \in \mathcal{C}^n(\mathbb{R}, X)$ with

$$\|Dy\|_\infty \leq \varepsilon \quad (3)$$

there exists $y_0 \in \ker D$ such that

$$\|y - y_0\|_\infty \leq K\varepsilon. \quad (4)$$

Denote by K_D the infimum of all constants K satisfying (4). If K_D is an Ulam constant of D , it is called the best Ulam constant or the Ulam constant of D . In this paper we are looking for the best Ulam constant of D .

EXPLICIT ALGEBRAIC SOLUTION TO ZOLOTAREV'S "FIRST PROBLEM" FOR POLYNOMIALS OF DEGREE $n \in \{6, 7, 8\}$

Heinz-Joachim Rack

*Dr. Rack Consulting GmbH, Steubenstrasse 26a, 58097 Hagen,
Germany*

[heinz-joachim.rack@drrack.com]

MSC 2010: 41A10, 41A29, 41A50

Keywords: Algebraic solution, approximation, explicit, first problem, least deviation from zero, polynomial, proper, Zolotarev.

For Zolotarev's so-called "First Problem", see [1], we provide an explicit algebraic solution for the polynomial degree $n \in \{6, 7, 8\}$. We have announced this result on a poster presented at the Poster Session of the IX Jaen Conference on Approximation Theory, see pp. 59-60 at URL <http://www.ujaen.es/revista/jja/jca/principal.pdf>

REFERENCES

- [1] B.C. Carlson, J. Todd, *Zolotarev's first problem – the best approximation by polynomials of degree $\leq n - 2$ to $x^n - n\sigma x^{n-1}$ in $[-1, 1]$* , Aequationes Math. 26 (1983), pp. 1-33.

APPROXIMATION BY MODIFIED U_n^ρ OPERATORS

Voichița Adriana Radu

Department of Statistics-Forecasts-Mathematics, Babeş-Bolyai University, Cluj-Napoca, Romania

[voichita.radu@econ.ubbcluj.ro]

MSC 2010: 41A10, 41A25, 41A36

Keywords: λ -Bernstein operators, U_n^ρ operators, moduli of continuity, rate of convergence, Voronovskaya type theorem.

The U_n^ρ operators was introduced in 2007 by R. Păltănea and since then, many researchers extend the study of them in vary directions. In our paper we consider a generalization of these operators using a new Bezier bases $\tilde{b}_{n,k}(\lambda, x)$ with shape parameter λ . Some approximation properties are given, including local approximation, error estimation in terms of moduli of continuity and Voronovskaja-type asymptotic formulas.

REFERENCES

- [1] A.M. Acu, H.H. Gonska, I. Raşa, *Gruss-type and Ostrowski-type inequalities in approximation theory*, Ukrainian Math. J., 63(6) (2011), pp. 843-864.
- [2] F. Altomare, M. Campiti, *Korovkin-type Approximation Theory and its Applications*, de Gruyter, Berlin, New York, 1994.
- [3] Q.B. Cai, B.Y. Lian, G. Zhou, *Approximation properties of λ -Bernstein operators*, Journal of Inequalities and Applications, 12 (2018), pp. 1-11.

- [4] I. Gavrea, H.H. Gonska, R. Păltănea, G. Tachev, *General estimates for the Ditzian-Totik modulus*, East J. Approx. Th., 9 (2003), pp. 175-194.
- [5] H.H. Gonska, G. Tachev, *Gruss-type inequalities for positive linear operators with second order moduli*, Matematicki Vesnik, 63(4), 2011, pp. 247-252.
- [6] R. Păltănea, *Approximation Theory Using positive Linear Operators*, Birkhauser, Boston, 2004.
- [7] R. Păltănea, *A class of Durrmeyer type operators preserving linear functions*, Ann. Tiberiu Popoviciu Sem. Funct. Equat. Approxim. Convex., Cluj-Napoca,5, 2007, pp. 109-117.
- [8] V.A. Radu, *Quantitative Estimates for Some Modified Bernstein-Stancu Operators*, Miskolc Math. Notes 19(1), 2018, pp. 517-525.

BOUNDS AND CLOSED FORMS FOR SPECIAL FUNCTIONS

Ioan Raşa

Department of Mathematics, Technical University of Cluj-Napoca, 28 Memorandumului Street, 400114 Cluj-Napoca, Romania

[ioan.rasa@math.utcluj.ro]

MSC 2010: 33E30, 33C05, 94A17

Keywords: Heun function, entropy, hypergeometric function.

We obtain new bounds and closed forms for some hypergeometric/Heun functions. The results are related to those presented in [1] and [2]. We also recall the following

CONJECTURE (I. Raşa, University of Bielsko-Biala, November 15, 2017)

$f \in C[0, 1]$ log-concave \Rightarrow

$$\sum_{\substack{i+j=h \\ 0 \leq i \leq n-1 \\ 0 \leq j \leq n}} \binom{n-1}{i} \binom{n}{j} \left((n-1-i) f\left(\frac{j}{n}\right) \Delta_{\frac{2}{n}}^2 f\left(\frac{i}{n}\right) - (n-j) \Delta_{\frac{1}{n}}^1 f\left(\frac{i}{n}\right) \Delta_{\frac{1}{n}}^1 f\left(\frac{j}{n}\right) \right) \leq 0,$$

for all $n \geq 1$, $h \in \{0, 1, \dots, 2n-2\}$.

REFERENCES

- [1] A. Bărar, G. Mocanu, I. Raşa, *Heun functions related to entropies*, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas. <https://doi.org/10.1007/s13398-018-0516-x>
- [2] A. Bărar, G. Mocanu, I. Raşa, *Bounds for some entropies and special functions*, Carpathian J. Math. 34 (2018), No. 1, pp. 9-15.

BETTER CONVERGENCE PROPERTIES OF CERTAIN POSITIVE LINEAR OPERATORS

Augusta Rațiu

Department of Mathematics and Computer Science, Lucian Blaga University, Sibiu, Romania
[augu2003@yahoo.com]

MSC 2010: 41A10, 41A25, 41A36.

Keywords: Linear positive operators, Bernstein operators.

In this paper there are presented the modified positive linear operators which depend on a certain function ρ defined on $[0, 1]$, and they have a better degree of approximation than classical ones. To illustrate this, some examples are given for Bernstein operators, Lupas operators and genuine Bernstein-Durrmeyer operators.

REFERENCES

- [1] T. Acar, A. Aral, I. Raşa, *Modified Bernstein-Durrmeyer operators*, General Mathematics, 22(1) (2014), pp. 27-41.
- [2] A.M. Acu, P.N. Agrawal, T. Neer, *Approximation properties of the modified Stancu operators*, Numerical Functional Analysis and Optimization, 38(3) (2017), pp. 279-292.
- [3] A.M. Acu, H. Gonska, I. Rasa, *Grüss-type and Ostrowski-type inequalities in approximation theory*, Ukrainian Math. J., 63(6) (2011), pp. 843-864.
- [4] O. Agratini, *Properties of discrete non-multiplicative operators*, Anal. Math. Phys., (2017), doi.org/10.1007/s13324-017-0186-4.

NUMERICAL OPTIMAL CONTROL FOR SATELLITE ATTITUDE PROFILES

Ralf Rigger

*Department Mathematik, Naturwissenschaften und
Datenverarbeitung, Technische Hochschule Mittelhessen,
Wilhelm-Leuschner-Strasse 13, 61169 Friedberg, Germany
[ralf.rigger@mnd.thm.de]*

MSC 2010: 49J15, 90C20, 90C30

Keywords: Satellite attitude profile, slew, optimal control, numerical simulation, NMPC.

Many modern science satellites are 3-axis stabilized. Attitude profiles play a central role in satellite control. Besides the dynamical properties numerous constraints need to be fulfilled. These additional constraints are usually of elementary geometric nature and can often be described as affine-linear functions. Several examples of problems of this type are stated and solutions with different optimal control methods [BH81] and respective numerical simulations will be presented. The generation of valid attitudes has often to consider the alignment of the solar arrays with sun line. In [RYMC04] a generic way for calculating such attitudes is given. By regarding the problem from an 'euler angle point of view', the remaining two degrees of freedom form a flat torus and allow the search for attitudes that obey further geometric constraints. Another way to approach this problem is to use optimal control methods. A method based on a simple cost function is proposed. Further some analytic background for understanding the choice of the cost function is given. Besides the attitude profile also the dynamics

has to be considered. Using numerical model predictive control [W09], [D15] an approximative filter to reduce spacecraft rates to a given level and the corresponding numerical results are presented.

REFERENCES

- [KHS96] K.-H. Spindler, *Optimal attitude control of a rigid body*, Applied Mathematics and Optimization, Volume 34, Number 1, 1996
- [RYMC04] R. Rigger, J. Yde, M. Mller, V. Companys, *The Optimisation of Attitude profiles for SMART-1: A highly constrained Problem*, Proceedings of the 18th International Symposium on Space Flight Dynamics, 2004
- [W09] L. Wang, *Model Predictive Control System Design and Implementation*, Springer - Advances in Industrial Control, 2009
- [D15] M. Diehl, *Spring School on Nonlinear Model Predictive Control*, 2015
- [BH81] A. E. Bryson & Y. C. Ho, *Applied Optimal Control: Optimization, Estimation & Control: Optimization Estimation and Control*, 1981
- [WK03] R. Wisniewski and P. Kulczycki, *Slew Maneuver Control for Spacecraft Equipped with Star Camera and Reaction Wheels*, Preprint submitted to Elsevier Science, 2003

MASTER SLAVE SYNCHRONIZATION FOR STOCHASTIC LATTICE SYSTEMS

Björn Schmalfuss

*Department of Mathematics, Friedrich Schiller University of
Jena, Germany*

[bjorn.schmalfuss@uni-jena.de]

MSC 2010: 60H15

Keywords: Random dynamical system, random inertial manifold, synchronization.

We consider two stochastic lattice systems. The nodes of each of these systems are linearly coupled. In addition there is a coupling for any node between these systems. We state the existence of an invariant random manifold in the phase space such that we have synchronization in the following sense. Any solution trajectory of the systems converges exponentially fast to some solution trajectory "living" on the random manifold.

APPROXIMATION WITH LOCAL ITERATED FUNCTION SYSTEMS

Anna Soós, Ildikó Somogyi*

Department of Mathematics, Faculty of Mathematics and Computer Science, Babeş-Bolyai University, Cluj-Napoca, Romania

MSC 2010: 65D05

Keywords: Fractal functions, attractors, interpolation.

The set of real data can be approximated with fractal interpolation functions. These fractal interpolation functions can be constructed with the so-called iterated function systems. Local iterated function systems are important generalization of iterated function systems. In order to obtain new approximation methods we can combine this methods with the classical interpolation methods. In this article using the fact that graphs of piecewise polynomial functions can be written as the fixed points of local iterated function system we give some delimitation of the error for this new approximation methods.

REFERENCES

- [1] M.F. Barnsley, *Fractals Everywhere*, Academic Press (1993)
- [2] M.F. Barnsley, M. Hegeland and P. Massopust, *Numerics and Fractals*, <https://arxiv.org/abs/1309.0972>, (2014)

- [3] J. Kolumbán, A. Soós, *Fractal functions using contraction method in probabilistic metric spaces*, Proceeding of the 7th International Multidisciplinary Conference Fractal 2002, Complexity and Nature, Emergent Nature, M.M. Novak (ed.), World Scientific 2002, pp. 255-265.
- [4] M.A. Nevascués, M.V. Sebastián, *Generalization of Hermite functions by fractal interpolation*, J. Approx. Theory, 131 (1) 2004, pp. 19-29.
- [5] H.Y. Wang, J.S. Yu, *Fractal interpolation functions with variable parameters and their analytical properties*, J. Approx Theory, 175 (2013), pp. 1-18.

STONE-WEIERSTRASS THEOREMS FOR RANDOM FUNCTIONS

Hans-Jörg Starkloff

Faculty of Mathematics and Computer Science, University of Technology, Freiberg, Saxonia, Germany

[Hans-Joerg.Starkloff@math.tu-freiberg.de]

MSC 2010: 41A10, 41A65, 60G99

Keywords: Random function, Stone-Weierstrass theorem, stochastic convergence.

Due to an increased use of random models and an intensive investigation of random ordinary and partial differential equations a more systematic treatment of approximation methods for random functions and random variables in function spaces seems to be appropriate. Basic results in deterministic approximation theory are related to different versions of Stone-Weierstrass theorems. In the talk some stochastic versions of Stone-Weierstrass theorems are presented, generalizing corresponding results of, e.g., [1] and [2].

REFERENCES

- [1] D. Dugué, *Traité de statistique théorique et appliquée*, Paris, 1958.
- [2] O. Onicescu, V.I. Istrăţescu, *Approximation theorems for random functions* Rendiconti di Matematica, serie vi, 8 (1975), pp. 65-81.

A HEURISTIC INTRODUCTION TO NUMERICAL MODELING OF DIFFUSION PROCESSES

Nicolae Suciu

*Tiberiu Popoviciu Institute of Numerical Analysis, Romanian
Academy, Cluj-Napoca, Romania*

[nsuciu@ictp.acad.ro]

MSC 2010: 60

Keywords: Diffusion, independent identically distributed (i.i.d.) random variables.

Sums of sequences of independent identically distributed (i.i.d.) random variables can be formulated by the Law of Large Numbers or the Central Limit Theorem as diffusion processes. This property of the i.i.d. random variables, which lies at the core of numerical modeling of diffusion processes, will be illustrated by generating various diffusion processes and further used to model diffusion at both microscopic and macroscopic level by numerical solutions of Ito and Fokker-Planck equations, respectively.

REFERENCES

- [1] C. Vamoş, N. Suciu and H. Vereecken, *Generalized random walk algorithm for the numerical modeling of complex diffusion processes*, J. Comp. Phys., 186 (2003), 527-544, doi: 10.1016/S0021-9991(03)00073-1

QUANTITATIVE ESTIMATES FOR q -BERNSTEIN OPERATORS IN MOBILE INTERVALS

E. Taş^{1,*}, T. Yurdakadim² and I. Sakaoglu³

¹*Department of Mathematics, Ahi Evran University, Turkey*

²*Department of Mathematics, Hitit University, Turkey*

³*Department of Mathematics, Ankara University, Turkey*

[emretas86@hotmail.com],

[tugbayurdakadim@hotmail.com],

[i.sakaoglu@gmail.com]

MSC 2010: 41A25, 41A36

Keywords: q -Bernstein operators, mobile interval.

In this paper we introduce q -Bernstein type operators which is an extension of classical case since they convert q -Bernstein operators if n is sufficiently large. We also study some approximation properties and obtain direct and inverse theorems.

This work was supported by the Ahi Evran University Scientific Research Projects Coordination Unit. Project Number: FEF.A4.18.004

REFERENCES

- [1] H.S. Jung, N. Deo, M. Dhamija, *Pointwise approximation by Bernstein type operators in mobile interval*, Appl. Math. Comput., 244 (2014), pp. 683-694.
- [2] V. Kac and P. Cheung, *Quantum Calculus*, springer-Verlag, 2002.

AN APPLICATION OF INVERSE PADÉ INTERPOLATION

Radu T. Trîmbițaș

*Department of Mathematics, “Babeș-Bolyai” University,
Cluj-Napoca, Romania*

[tradu@math.ubbcluj.ro]

MSC 2010: 65H05,65Y99, 65-0

Keywords: Nonlinear equations, Padé approximation, Maple.

We use inverse Padé interpolation and Maple to find the following fourth order method for the solution of a scalar nonlinear equation

$$\Phi(x) = x - \frac{f(x)}{f'(x)} \left\{ 1 + \frac{\frac{1}{2}}{\frac{f'(x)}{f''(x)} \left[\frac{f'(x)}{f(x)} + \frac{f'''(x)}{3f'(x)} \right] - 1} \right\}.$$

Also a result on convergence and a MATLAB implementation are given.

REFERENCES

- [1] W. Gander, M. J. Gander, F. Kwok, *Scientific Computing. An introduction using Maple and MATLAB*, Springer, 2014.

SOME APPROXIMATION PROPERTIES OF GENERALIZED MEYER-KÖNIG AND ZELLER OPERATORS VIA ABEL CONVERGENCE

Mehmet Ünver^{1,*} and Dilek Söylemez²

¹*Department of Mathematics, Faculty of Science, Ankara University, Turkey*

²*Department of Computer Programming, Elmadağ Vocational School, Ankara University, Turkey*

[munver@ankara.edu.tr], [dsoylemez@ankara.edu.tr]

MSC 2010: 41A36, 40A35, 40G10

Keywords: Abel convergence, Korovkin type theorem, modulus of continuity.

Korovkin type approximation theory mainly deals with the convergence of a sequence of positive linear operators defined on the space of real-valued continuous functions on a closed interval. If the ordinary convergence does not work then we may consider a summability method that is consistent with and stronger than ordinary convergence. In this talk, using the Abel convergence method we study some approximation properties of generalized Meyer-König and Zeller operators. Moreover, we obtain the rate of convergence by means of modulus of continuity.

REFERENCES

- [1] Ö. Atlıhan, M. Ünver, *Abel transforms of convolution operators*, Georgian Math. J. 22 (3), (2015) pp. 323-329.
- [2] J. Boos, *Classical and Modern Methods in Summability*, Oxford, UK, Oxford University Press, (2000).
- [3] O. Duman, M.K. Khan, C. Orhan, *A-Statistical convergence of approximating operators*, Math Inequal. Appl. 6, (2003), 689-699. 90, (1953), pp. 961–964.
- [4] A.D. Gadjiev, C. Orhan, *Some approximation theorems via statistical convergence*, Rocky Mountain J. Math. 32, (2002), pp. 129–138.
- [5] P.P. Korovkin, *Convergence of linear positive operators in the spaces of continuous functions* (Russian), Doklady Akad. Nauk. SSSR (N.S)
- [6] M.W. Mller, *Ueber die Ordnung der Approximation durch die Folge der Operatoren von Meyer-Knig und Zeller und durch die Folge deren erster Ableitungen*, Bul. Inst. Politechn. Iasi, 14 (1968), pp. 83-90.
- [7] R.E. Powell, S.M. Shah, *Summability Theory and Its Applications*, Prentice-Hall of India, New Delhi, (1988).
- [8] İ. Sakaoglu, M. Ünver, *Statistical approximation for multi-variable integrable functions*, Miskolc Math. Notes 13, no. 2, (2012), pp. 485-491.
- [9] M. Ünver, *Abel transforms of positive linear operators*, ICNAAM 2013, AIP Conference Proceedings, Volume 1558, pp. (2013), pp. 1148-1151.

POSSIBLY INFINITE GENERALIZED ITERATED FUNCTION SYSTEMS COMPRISING φ -MAX CONTRACTIONS

Silviu Urziceanu

*Department of Mathematics and Computer Science,
University of Pitești, Romania, Târgul din Vale 1, 110040,
Pitești, Argeș, Romania*
[fmi_silviu@yahoo.com]

MSC 2010: 28A80, 37C70, 41A65, 54H25

Keywords: Possibly infinite generalized iterated function system,
 φ -max contraction, attractor, canonical projection.

One way to generalize the concept of iterated function system was proposed by R. Miculescu and A. Mihail (see [3] and [4]) under the name of generalized iterated function system (for short GIFS). More precisely, given $m \in \mathbb{N}^*$ and a metric space (X, d) , a generalized iterated function system of order m is a finite family of functions $f_1, \dots, f_n : X^m \rightarrow X$ satisfying certain contractive conditions. Another generalization of the notion of iterated function system is given by those systems consisting of φ -max contractions (see [1]).

Combining these two lines of research, we prove that the fractal operator associated to a possibly infinite generalized iterated function system comprising φ -max contractions is a Picard operator (whose fixed point is called the attractor of the system). Paper [2] inspired us to associate to each possibly infinite generalized iterated function system comprising φ -max contractions \mathcal{F} (of order m) an operator

$H_{\mathcal{F}} : \mathcal{C}^m \rightarrow \mathcal{C}$, where \mathcal{C} stands for the space of continuous and bounded functions from the shift space on the metric space corresponding to the system. We prove that $H_{\mathcal{F}}$ is a Picard operator whose fixed point is the canonical projection associated to \mathcal{F} .

REFERENCES

- [1] F. Georgescu, R. Miculescu, A. Mihail, *A study of the attractor of a φ -max-IFS via a relatively new method*, J. Fixed Point Theory Appl., (2018) 20:24, <https://doi.org/10.1007/s11784-018-0497-6>.
- [2] A. Mihail, *The canonical projection between the shift space of an IIFS and its attractor as a fixed point*, Fixed Point Theory Appl., 2015, Paper No. 75, 15 p.
- [3] A. Mihail, R. Miculescu, *Applications of Fixed Point Theorems in the Theory of Generalized IFS*, Fixed Point Theory, Appl. Volume 2008, Article ID 312876, 11 pages doi: 10.1155/2008/312876.
- [4] A. Mihail, R. Miculescu, *Generalized IFSs on Noncompact Spaces*, Fixed Point Theory Appl., Volume 2010, Article ID 584215, 11 pages doi: 10.1155/2010/584215.

APPROXIMATION BY BIVARIATE NONLINEAR SINGULAR INTEGRAL OPERATORS IN MOBILE RECTANGLE

Gümrah Uysal

*Department of Computer Technologies, Division of Technology
of Information Security, Karabük University, 78050, Karabük,
Turkey*

[e-mail:fgumrahuyosal@gmail.com]

MSC 2010: 41A35, 41A25, 47G10

Keywords: μ -generalized Lebesgue point, rate of convergence,
Lipschitz condition, nonlinear analysis.

Let $\Lambda \subset [0, \infty; 0, \infty]$ be a two-dimensional index set consisting of $\sigma = (\xi, \eta)$ indices with accumulation point $\sigma_0 = (\xi, \eta)$. We assume that the function $K_\sigma : \mathbb{R}^2 \times \mathbb{R} \rightarrow \mathbb{R}$ is Lebesgue integrable on \mathbb{R}^2 and satisfies some conditions including Lipschitz condition with respect to its third variable for any $\sigma \in \Lambda$. The main purpose of this work is to investigate the conditions under which Fatou type pointwise convergence is obtained for the operators in the following setting:

$$(T_\sigma f)(x, y) = \int_{a-\xi c-\eta}^{b+\xi d+\eta} \int K_\sigma(t-x, s-y, f(t, s)) ds dt, \quad (x, y) \in \mathbf{R},$$

where a, b, c and d are arbitrary real numbers and $\mathbf{R} = (a - \xi, b + \xi; c - \eta, d + \eta)$ is a mobile rectangle, at μ -generalized Lebesgue point of measurable function $f \in L_1^\omega(\mathbf{R})$,

as $(x, y, \sigma) \rightarrow (x_0, y_0, \sigma_0)$. Here, $L_1^\omega(\mathbf{R})$ is the space of all measurable functions f for which $\left|\frac{f}{\omega}\right|$ is integrable over \mathbf{R} , where $\omega : \mathbb{R}^2 \rightarrow \mathbb{R}^+$ is a weight function satisfying some extra conditions. The obtained results are used for presenting some theorems for the rate of convergences.

REFERENCES

- [1] A.D. Gadjiev, *The order of convergence of singular integrals which depend on two parameters*, Special problems of functional analysis and their applications to the theory of differential equations and the theory of functions. Izdat. Akad. Nauk Azerbaidžan. SSR., Baku (1968), pp. 40-44.
- [2] B. Rydzewska, *Approximation des fonctions de deux variables par des intégrales singulières doubles*, Fasc. Math., 8 (1974), pp. 35–45.
- [3] C. Bardaro, J. Musielak, G. Vinti, *Nonlinear Integral Operators and Applications*, De Gruyter Series in Nonlinear Analysis and Applications, 9. Walter de Gruyter & Co., Berlin (2003).
- [4] G. Uysal, M.M. Yilmaz, E. Ibikli, *On pointwise convergence of bivariate nonlinear singular integral operators*, Kuwait J. Sci. 44 (2) (2017), pp. 46–57.
- [5] H.S. Jung, N. Deo, M. Dhamija, *Pointwise approximation by Bernstein type operators in mobile interval*, Appl. Math. Comput. 244 (2014), pp. 683–694.
- [6] H. Karsli, *Convergence and rate of convergence by nonlinear singular integral operators depending on two parameters*, Appl. Anal., 85 (6-7) (2006), 781–791.
- [7] J. Musielak, *On some approximation problems in modular spaces*, Constructive function theory, Proceedings of International Conference Varna, 1-5 June, 1981, Publication House of Bulgarian Academic of Sciences, Sofia, (1983), pp. 455-461.

- [8] M. Dhamija, R. Sakai, N. Deo, *On approximation by Phillips type modified Bernstein operator in a mobile interval*, J. Class. Anal. 7 (1) (2015), pp. 25–37.
- [9] R. Taberski, *Singular integrals depending on two parameters*, Prace Mat., 7 (1962), pp. 173–179.
- [10] R. Taberski, *On double integrals and Fourier series*, Ann. Polon. Math., 15 (1964), pp. 97–115.
- [11] R. Taberski, *On double singular integrals*, Comment. Math. Prace Mat., 19 (1) (1976), pp. 155-160.
- [12] S. Siudut, *On the convergence of double singular integrals*, Comment. Math. Prace Mat., 28 (1), (1988), pp. 143–146.
- [13] T. Swiderski, E. Wachnicki, *Nonlinear singular integrals depending on two parameters*, Comment. Math., 40 (2000), pp. 181-189.

EXPLICIT ALGEBRAIC SOLUTION OF THE ZOLOTAREV PROBLEM VIA GROEBNER BASIS

Robert Vajda

Bolyai Institute, University of Szeged, Hungary
[vajdar@math.u-szeged.hu]

MSC 2010: 41A10, 41A29, 41A50

Keywords: Differential equation, equioscillation, equiripple, explicit form, Groebner basis, least deviation from zero, Mathematica, polynomial, proper, symbolic computation, Zolotarev.

The exact general solution to Zolotarev's First Problem was known only in terms of elliptic integrals and functions [1]. Interests in giving explicit algebraic descriptions to small degree Zolotarev polynomials revived in some recent work [2, 3]. This talk investigates the power of symbolic computational tools (Groebner basis and quantifier elimination) to solve the problem. It turns out that generic polynomial expressions of a symbolic parameter can be obtained via Groebner basis method using the defining differential equation of the proper Zolotarev polynomials at least up to degree eight. The knowledge of explicit algebraic solutions can be used for solving various related computational problems in approximation theory. Most of the computation is done with aid of the Wolfram Mathematica computer algebra system.

REFERENCES

- [1] B. C. Carlson, J. Todd, *Zolotarev's first problem - the best approximation by polynomials of degree $\leq n - 2$ to $x^n - n\sigma x^{n-1}$ in $[-1, 1]$* , Aequationes Math. 26 (1983), pp. 1-33.
- [2] V. A. Malyshev, *Algebraic solution of the Zolotarev problem*, Algebra i Analiz, 14(4) (2002), pp. 238-240.
- [3] H.-J. Rack, *Explicit algebraic solution to Zolotarev's First Problem for polynomials of degree $n \in \{6, 7, 8\}$* , IX Jaen Conference on Approximation Theory (2018), Poster session.

FINITE DIFFERENCE SCHEME FOR THE HIGH ORDER NONLINEAR SCHRÖDINGER EQUATION WITH LOCALIZED DISSIPATION.

Rodrigo Véjar Asem

Centro de Investigación en Ingeniería Matemática (CI²MA)
Departamento de Ingeniería Matemática (DIM)
Universidad de Concepción, Chile
[rodrigovejar@rocketmail.com,
rodrigovejar@ing-mat.udec.cl]

Keywords: HNLS, damping, dissipation, finite difference method.

A Finite Difference scheme for the High Order Nonlinear Schrödinger (HNLS) equation in 1D, with localized damping, will be presented. The equation can model superluminal optical solitons in atomic systems, as well as femtosecond solitons travelling in optical fibers made by different materials. The method can preserve the numerical energy, and can almost preserve the numerical charge. Numerical results will be shown.

This is a joint work with Marcelo M. Cavalcanti (UEM-Brazil), Wellington J. Corrêa (UTFPR-Brazil), and Mauricio Sepúlveda C. (CI²MA - DIM, UdeC, Chile).

REFERENCES

- [1] M. Delfour, M. Fortin, G. Payre, *Finite-Difference Solutions of a Non-linear Schrödinger Equation*. Journal of Computational Physics **44** (1981), pp. 277-288.
- [2] D. Furihata, T. Matsuo, *Discrete variational derivative method: a structure-preserving numerical method for partial differential equations*, Chapman and Hall, ISBN 978-1-4200-9445-9 (2011).
- [3] Y. Kodama, A. Hasegawa, *Nonlinear Pulse Propagation in a Monomode Dielectric Guide*. IEEE Journal of Quantum Electronics, **QE-23** (1987), pp. 510-524.
- [4] T.L. Perel'man, A. Kh. Fridman, M.M. El'yashevich. *A modified Korteweg-de Vries equation in electrohydrodynamics*. Sov. Phys. JETP, **39**, No. 4 (1974).
- [5] M.J. Potasek, M. Tabor., *Exact solutions for an extended nonlinear Schrödinger equation*. Physics Letters A, Volume 154 (1991), number 9.
- [6] V.M. Vyas et al., *Chirped chiral solitons in the nonlinear Schrödinger equation with self-steepening and self-frequency shift*. Physical Review A, **78** (2008), 021803(R).

NON-ALMOST PERIODIC SOLUTIONS OF DIFFERENCE SYSTEMS VIA ITERATIVE METHODS

Michal Veselý

*Department of Mathematics and Statistics, Faculty of Science,
Masaryk University, Czech Republic*

[michal.vesely@mail.muni.cz]

MSC 2010: 12H10, 39A06, 39A24, 42A75, 43A60

Keywords: Limit periodicity, almost periodicity, limit periodic sequences, asymptotically almost periodic solutions, difference equations.

This talk is based on the joint works [1, 2, 3] with P. Hasil. Using iterative methods, we study the solution spaces of limit periodic homogeneous linear difference systems, where the coefficient matrices of the considered systems are taken from a given commutative group which does not need to be bounded. In particular, we study such systems whose fundamental matrices are not asymptotically almost periodic or which have solutions vanishing at infinity. We identify conditions on the matrix group which guarantee that these systems form a dense subset in the space of all considered systems. Note that the elements of the coefficient matrices are taken from an infinite field with an absolute value and that the corresponding almost periodic case is treated as well.

REFERENCES

- [1] P. Hasil, M. Veselý, *Limit periodic homogeneous linear difference systems*, Appl. Math. Comput., 265 (2015), pp. 958–972.
- [2] P. Hasil, M. Veselý, *Solution spaces of homogeneous linear difference systems with coefficient matrices from commutative groups*, J. Difference Equ. Appl., 23 (2017), no. 8, pp. 1324–1353.
- [3] P. Hasil, M. Veselý, *Asymptotically almost periodic solutions of limit periodic difference systems with coefficients from commutative groups*, submitted (2018).

OPTIMIZATION ALGORITHMS BASED ON OPERATOR SPLITTING AND ENERGY-PRESERVING NUMERICAL INTEGRATORS

Adrian Viorel^{1,*} and Cristian Daniel Alecsa²

¹ *Department of Mathematics, Technical University of Cluj-Napoca, Romania*

² *Tiberiu Popoviciu Institute of Numerical Analysis, Romanian Academy, Romania*

[Adrian.Viorel@math.utcluj.ro],

[cristian.alecsa@ictp.acad.ro]

MSC 2010: 65J08, 90C59, 90C30

Keywords: Optimization algorithm, dynamical system, Strang splitting, conservative numerical method.

The present contribution deals with nonconvex optimization from a dynamical systems perspective. More precisely, we consider the second order evolution equation

$$\ddot{u} + \dot{u} = -\nabla G(u) \tag{1}$$

which can be understood (cf. [1]) as describing a damped mechanical system with the nonconvex objective function $G : \mathbb{R}^n \rightarrow \mathbb{R}$ as the system's potential energy. Its asymptotic behavior has been analyzed quite recently by Bégout et. al. who show the convergence of solutions, at precise rates, to minimizers of G .

A similar approach applies to the vanishing damping model

$$\ddot{u} + \frac{\gamma}{t}\dot{u} = -\nabla G(u), \quad (2)$$

which is the continuous counterpart of the celebrated Nesterov optimization algorithm (cf. [5]).

In this context, our aim is to investigate discrete dynamical systems which faithfully reproduce the properties of (1) or (2). These discrete dynamical systems are obtained by means of a two-stage discretization of (1) or (2), consisting in a Strang splitting semidiscretization that separates the conservative and dissipative parts of the system followed by an energy-preserving numerical integration of the conservative part.

The Strang splitting approximation (see [2] and references therein)

$$e^{h(A+B)} \approx e^{\frac{h}{2}A} e^{hB} e^{\frac{h}{2}A}, \quad e^{t(A+B)} \approx \left(e^{\frac{h}{2}A} e^{hB} e^{\frac{h}{2}A} \right)^n, \quad t = nh$$

is known to have an improved order of approximation when compared to the simpler Lie splitting

$$e^{h(A+B)} \approx e^{hA} e^{hB}, \quad e^{t(A+B)} \approx \left(e^{hA} e^{hB} \right)^n, \quad t = nh,$$

as both are dealing with the linear problem $\dot{z} = Az + Bz$ by iteratively solving two subproblems: $\dot{z} = Az$ and $\dot{z} = Bz$. Recent progress [3], however, has shown that the semilinear problem $\dot{z} = Az + f(z)$ can be treated in exactly the same manner by only replacing e^{hB} with the solution operator (nonlinear semigroup) $\mathcal{U}(h)$ of $\dot{z} = f(z)$.

The evolution equations (1), (2) both fit nicely in this semilinear framework if rewritten as systems of first order equations

$$\begin{cases} \dot{u} = v, \\ \dot{v} = -c(t)v - \nabla G(u), \end{cases} \quad (3)$$

with either $c(t) = 1$ or $c(t) = \gamma/t$. Furthermore, the splitting method can be applied in such a way that the conservative and dissipative parts of the system are separated, the subproblems that need to be solved iteratively being

$$\text{(Cons)} \begin{cases} \dot{u} = v, \\ \dot{v} = -\nabla G(u) \end{cases} \quad \text{and} \quad \text{(Dissip)} \begin{cases} \dot{u} = 0, \\ \dot{v} = -c(t)v. \end{cases}$$

The resulting semidiscretization generates a discrete dynamical system which has

- (i) the same equilibria $(u^*, 0)$, with $\nabla G(u^*) = 0$, as (3);
- (ii) the same Lyapunov function $H(u, v) = \frac{1}{2} \|v\|^2 + G(u)$ as (3);
- (iii) an $O(h^2)$ accuracy on finite time intervals.

While the dissipative subproblem is explicitly solvable, the conservative one can be integrated using an energy-preserving numerical scheme (see [4]) such that all the properties (i), (ii) and (iii) remain valid.

Further, the infinite-dimensional case is also discussed.

REFERENCES

- [1] H. Attouch, X. Goudou, and P. Redont, *The heavy ball with friction method. I. The continuous dynamical system: global exploration of the local minima of a real-valued function by asymptotic analysis of a dissipative dynamical system*, Commun. Contemp. Math., 2(1), 2000, pp. 1–34.
- [2] R. Glowinski and S.J. Osher, editors, *Splitting methods in communication, imaging, science, and engineering*, Scientific Computation. Springer, Cham, 2016.
- [3] E. Hansen, F. Kramer, and A. Ostermann, *A second-order positivity preserving scheme for semilinear parabolic problems*, Appl. Numer. Math., 62(10), 2012, pp. 1428–1435.
- [4] G.R.W. Quispel and D.I. McLaren, *A new class of energy-preserving numerical integration methods*, J. Phys. A, 41(4):045206, 7, 2008.
- [5] W. Su, S. Boyd, and E.J. Candès, *A differential equation for modeling Nesterov’s accelerated gradient method: theory and insights*, J. Mach. Learn. Res., 17:Paper No. 153, 43, 2016.

DUAL BERNSTEIN POLYNOMIALS: NEW PROPERTIES AND FAST COMPUTATION

Filip Chudy¹, Pawel Woźny^{2,*}

²*Institute of Computer Science, University of Wrocław,
ul. Joliot-Curie 15, 50-383 Wrocław, Poland*

[pwo@cs.uni.wroc.pl]

MSC 2010: 33C45, 65Q30

Keywords: Differential equations, recurrence relations, Bernstein basis polynomials, dual Bernstein polynomials, Jacobi polynomials, Hahn polynomials.

Dual Bernstein polynomials introduced in [2], which are associated with the shifted Jacobi inner product, have recently found many applications in numerical analysis and computer graphics (curve intersection using Bézier clipping, degree reduction and merging of Bézier curves, polynomial approximation of rational Bézier curves, numerical solving of boundary value problems, numerical solving of fractional partial differential equations, etc.). Note that skillful use of these polynomials often results in less costly algorithms which solve some computational problems.

In the talk, we give new differential-recurrence properties of dual Bernstein polynomials which follow from relations between dual Bernstein and orthogonal Hahn and Jacobi polynomials. Using these results, the fourth-order differential equation satisfied by dual Bernstein polynomials has been constructed. Next, we obtain a fourth-order recurrence relation for these polynomials. The latter result may be useful in fast evaluation of dual Bernstein polynomials and their linear combinations or integrals which are

related to the least-square approximation in Bézier form. See also [1].

REFERENCES

- [1] F. Chudy, P. Woźny, *Differential-recurrence properties of dual Bernstein polynomials*, Applied Mathematics and Computation 2018, *accepted* (<https://arxiv.org/abs/1803.01735>).
- [2] S. Lewanowicz, P. Woźny, *Dual generalized Bernstein basis*, Journal of Approximation Theory 138 (2006), pp. 129–150.

HIGH-FREQUENCY EXPERT OPINIONS AND POWER UTILITY MAXIMIZATION IN A MARKET WITH GAUSSIAN DRIFT

Ralf Wunderlich

*Brandenburg University of Technology Cottbus – Senftenberg,
Germany*

[ralf.wunderlich@b-tu.de]

We consider a continuous-time financial market with partial information on the drift and solve utility maximization problems which include expert opinions on the unobservable drift. Stock returns are driven by a Brownian motion and the drift depends on a factor process which is an Ornstein Uhlenbeck process. Thus the drift is hidden and has to be estimated from observable quantities. If the investor only observes stock prices then the best estimate is the Kalman filter.

However, to improve the estimate, an investor may also rely on expert opinions providing a noisy estimate of the current state of the drift. This reduces the variance of the filter and thus improves expected utility. That procedure can be seen as a continuous-time version of the classical Black-Litterman approach.

For the associated portfolio problem with logarithmic utility explicit solutions are available in the literature. In this talk we consider the case of power utility. Here, we apply dynamic programming techniques and solve the corresponding dynamic programming equation for the value function. In particular we investigate the asymptotic behavior of the filter for high-frequency experts and derive

limit theorems for two different asymptotic regimes. In the first, variances of the expert opinions grow linearly with the arrival frequency while in the second they are constant. The derived limit theorems allow for simplified approximate solutions of utility maximization problems since the convergence of the filter carries over to the convergence of the value function. Numerical results are presented.

The talk is based on joint work with A. Gabih, H. Kondakji, J. Sass and D. Westphal.

KOROVKIN TYPE APPROXIMATION THEOREMS IN WEIGHTED SPACES VIA POWER SERIES METHOD

T. Yurdakadim^{1,*}, E. Taş² and Ö. G. Atlihan³

¹*Department of Mathematics, Hitit University, Turkey*

²*Department of Mathematics, Ahi Evran University, Turkey*

³*Department of Mathematics, Pamukkale University, Turkey*

[tugbayurdakadim@hotmail.com],

[emretas86@hotmail.com],

[oatlihan@pau.edu.tr]

MSC 2010: 41A36, 41A25

Keywords: Power series method, Korovkin type theorem, weighted space, rates of convergence.

In this paper we consider power series method which is also member of the class of all continuous summability methods. We study a Korovkin type approximation theorem for a sequence of positive linear operators acting from a weighted space C_{ρ_1} into a weighted space B_{ρ_2} with the use of the power series method which includes both Abel and Borel methods. We also consider the rates of convergence of these operators.

This paper was supported by the department of Scientific Research Projects of Hitit University. Project No: 19008

REFERENCES

- [1] O.G. Atlihan, C. Orhan, *Matrix summability and positive linear operators*, Positivity, 11 (2007), pp. 387-398.

LIST OF SPEAKERS

Ulrich Abel

*Department Mathematik, Naturwissenschaften und
Datenverarbeitung, Technische Hochschule Mittelhessen,
Wilhelm-Leuschner-Strasse 13, 61169 Friedberg, Germany*
[ulrich.abel@mnd.thm.de]

Ana Maria Acu

*Department of Mathematics and Informatics, Lucian Blaga
University, Sibiu, Romania*
[anamaria.acu@ulbsibiu.ro]

Mohd. Ahasan

*Department of Mathematics, Aligarh Muslim University,
Aligarh, 202002, India*
[ahasan.amu@gmail.com]

Francesco Altomare

Department of Mathematics, University of Bari, Italy
[francesco.altomare@uniba.it]

Alina Baias

*Department of Mathematics, Technical University of
Cluj-Napoca, Romania*
[Baias.Alina@math.utcluj.ro]

Carlo Bardaro

*Department of Mathematics and Computer Sciences,
University of Perugia, Italy*
[carlo.bardaro@unipg.it]

Marius-Mihai Birou

*Department of Mathematics, Technical University of Cluj
Napoca, Romania*
[Marius.Birou@math.utcluj.ro]

Mirella Cappelletti Montano

Department of Mathematics, University of Bari, Italy
[mirella.cappellettimontano@uniba.it]

Emil Căținaș

*Tiberiu Popoviciu Institute of Numerical Analysis, Romanian
Academy, Cluj-Napoca, Romania*
[ecatinas@ictp.acad.ro]

Larisa Cheregi

*Department of Mathematics, Technical University of
Cluj-Napoca, Romania*
[larisa.cheregi@math.utcluj.ro]

Maria Crăciun

*Tiberiu Popoviciu Institute of Numerical Analysis, Romanian
Academy, Cluj-Napoca, Romania*
[craciun@ictp.acad.ro]

Markus Dietz

Faculty of Mathematics and Computer Science, Technische Universität Bergakademie Freiberg, Saxony, Germany

[Markus.Dietz@math.tu-freiberg.de]

Julian Dimitrov

Department of Mathematics, University of Mining and Geology, Sofia, Bulgaria

[juldim@abv.bg]

María J. Garrido-Atienza

Departamento de Ecuaciones Diferenciales y Análisis Numérico, Universidad de Sevilla, Spain

[mgarrido@us.es]

Sorin G. Gal

Department of Mathematics and Computer Science, University of Oradea, Romania

[galso@uoradea.ro]

Bogdan Gavrea

Department of Mathematics, Technical University of Cluj-Napoca, Romania

[bogdan.gavrea@math.utcluj.ro]

Ioan Gavrea

Department of Mathematics, Technical University, Cluj-Napoca, Romania

[ioan.gavrea@math.utcluj.ro]

Flavian Georgescu

*Department of Mathematics and Computer Science,
University of Pitești, Romania, Pitești, Argeș, Romania*
[faviu@yahoo.com]

Călin-Ioan Gheorghiu

*Romanian Academy, "T. Popoviciu" Institute of Numerical
Analysis, Cluj-Napoca, Romania*
[ghcalin@ictp.acad.ro]

Heiner Gonska

University of Duisburg - Essen, Germany
[heiner.gonska@uni-due.de]

Wilfried Grecksch

*Institute of Mathematics, Martin-Luther-University of
Halle-Wittenberg, Germany*
[wilfried.grecksch@mathematik.uni-halle.de]

Petr Hasil

*Department of Mathematics and Statistics, Faculty of Science,
Masaryk University, Czech Republic*
[hasil@mail.muni.cz]

Margareta Heilmann

*School of Mathematics and Natural Sciences, University of
Wuppertal, Gaußstraße 20, D-42119 Wuppertal, Germany*
[heilmann@math.uni-wuppertal.de]

Harun Karsli

*Department of Mathematics, University of Bolu Abant Izzet
Baysal, Turkey*

[karsli_h@ibu.edu.tr]

Davod Khojasteh Salkuyeh

Department of Mathematics, University of Guilan, Rasht, Iran

[khojasteh@guilan.ac.ir]

Szilárd Csaba László

*Department of Mathematics, Technical University of
Cluj-Napoca, Romania*

[laszlosziszi@yahoo.com]

Vita Leonessa

*Department of Mathematics, Computer Science and
Economics, University of Basilicata, Potenza, Italy*

[vita.leonessa@unibas.it]

Aaron Melman

*Department of Applied Mathematics, Santa Clara University,
CA, USA*

[amelman@scu.edu]

Dan Miclăuș

*Department of Mathematics and Computer Science, Technical
University of Cluj-Napoca, North University Center at Baia
Mare, Romania*

[dan.miclaus@cunbm.utcluj.ro]

Radu Miculescu

*Department of Mathematics and Computer Science,
University of Bucharest, Romania, Academiei Street 14,
010014, Bucharest, Romania*
[miculesc@yahoo.com]

Gradimir V. Milovanović

*Mathematical Institute of the Serbian Academy of Sciences
and Arts, Serbia*
[gvm@mi.sanu.ac.rs]

Nicușor Minculete

Transilvania University, Romania
[minculetenu@yahoo.com]

Alexandru Mitrea

*Department of Mathematics, Technical University of
Cluj-Napoca, Romania*
[alexandru.ioan.mitrea@math.utcluj.ro]

Carmen Violeta Muraru

*Department of Mathematics and Informatics, Vasile
Alecsandri University, Bacau, Romania*
[cmuraru@ub.ro]

Vicuta Neagos

*Department of Mathematics, Technical University of
Cluj-Napoca, Romania*
[vicuta.neagos@math.utcluj.ro]

Adonia-Augustina Opreș

*Department of Mathematics, Technical University of
Cluj-Napoca, str. Memorandumului, nr. 28, 400144,
Cluj-Napoca, Romania*
[mate.salaj@yahoo.com]

Cihan Orhan

University of Ankara, Turkey
[Cihan.Orhan@science.ankara.edu.tr]

Diana Otrocol

*Department of Mathematics, Technical University of
Cluj-Napoca, 28 Memorandumului Street, 400114 Cluj-Napoca,
Romania*
[Diana.Otrocol@math.utcluj.ro]

Radu Păltănea

*Department of Mathematics and Informatics, "Transilvania"
University of Brașov, Romania*
[radupaltanea@unitbv.ro]

Flavius Pătrulescu

*Tiberiu Popoviciu Institute of Numerical Analysis, Romanian
Academy, Cluj-Napoca, Romania*
[fpatrulescu@ictp.acad.ro]

Daniel N. Pop

*Faculty of Engineering, Lucian Blaga University, Sibiu,
Romania*
[nicolae.pop@ulbsibiu.ro,popdaniel31@yahoo.com]

Dorian Popa

*Department of Mathematics, Technical University of
Cluj-Napoca, Romania*
[Popa.Dorian@math.utcluj.ro]

Heinz-Joachim Rack

*Dr. Rack Consulting GmbH, Steubenstrasse 26a, 58097 Hagen,
Germany*
[heinz-joachim.rack@drrack.com]

Voichița Adriana Radu

*Department of Statistics-Forecasts-Mathematics, Babeș-Bolyai
University, Cluj-Napoca, Romania*
[voichita.radu@econ.ubbcluj.ro]

Ioan Rașa

*Department of Mathematics, Technical University of
Cluj-Napoca, 28 Memorandumului Street, 400114 Cluj-Napoca,
Romania*
[ioan.rasa@math.utcluj.ro]

Augusta Rațiu

*Department of Mathematics and Computer Science, Lucian
Blaga University, Sibiu, Romania*
[augu2003@yahoo.com]

Ralf Rigger

*Department Mathematik, Naturwissenschaften und
Datenverarbeitung, Technische Hochschule Mittelhessen,
Wilhelm-Leuschner-Strasse 13, 61169 Friedberg, Germany*
[ralf.rigger@mnd.thm.de]

Björn Schmalfuss

*Department of Mathematics, Friedrich Schiller University of
Jena, Germany*

[bjourn.schmalfuss@uni-jena.de]

Ildikó Somogyi

*Department of Mathematics, Faculty of Mathematics and
Computer Science, Babeş-Bolyai University, Cluj-Napoca,
Romania*

Hans-Jörg Starkloff

*Faculty of Mathematics and Computer Science, University of
Technology, Freiberg, Saxonia, Germany*

[Hans-Joerg.Starkloff@math.tu-freiberg.de]

Nicolae Suci

*Tiberiu Popoviciu Institute of Numerical Analysis, Romanian
Academy, Cluj-Napoca, Romania*

[nsuciu@ictp.acad.ro]

E. Taş

Department of Mathematics, Ahi Evran University, Turkey

[emretas86@hotmail.com]

Radu T. Trîmbiţaş

*Department of Mathematics, Babeş-Bolyai University,
Cluj-Napoca, Romania*

[tradu@math.ubbcluj.ro]

Mehmet Ünver

*Department of Mathematics, Faculty of Science, Ankara
University, Turkey*

[munver@ankara.edu.tr]

Silviu Urziceanu

*Department of Mathematics and Computer Science,
University of Pitești, Romania, Târgul din Vale 1, 110040,
Pitești, Argeș, Romania*
[fmi_silviu@yahoo.com]

Gümrah Uysal

*Department of Computer Technologies, Division of Technology
of Information Security, Karabük University, 78050, Karabük,
Turkey*
[e-mail:fgumrahuyosal@gmail.com]

Robert Vajda

Bolyai Institute, University of Szeged, Hungary
[vajdar@math.u-szeged.hu]

Rodrigo Véjar Asem

*Centro de Investigación en Ingeniería Matemática (CI²MA)
Departamento de Ingeniería Matemática (DIM)
Universidad de Concepción, Chile*
[rodrigovejar@rocketmail.com,rodrigovejar@ing-mat.udec.cl]

Michal Veselý

*Department of Mathematics and Statistics, Faculty of Science,
Masaryk University, Czech Republic*
[michal.vesely@mail.muni.cz]

Adrian Viorel

*Department of Mathematics, Technical University of
Cluj-Napoca, Romania*
[Adrian.Viorel@math.utcluj.ro]

Wolfgang. L. Wendland

IANS & Simtech University Stuttgart, Germany
[wendland@mathematik.uni-stuttgart.de]

Pawel Woźny

*Institute of Computer Science, University of Wrocław,
ul. Joliot-Curie 15, 50-383 Wrocław, Poland*
[pwo@cs.uni.wroc.pl]

Ralf Wunderlich

*Brandenburg University of Technology Cottbus – Senftenberg,
Germany*
[ralf.wunderlich@b-tu.de]

T. Yurdakadim

Department of Mathematics, Hitit University, Turkey
[tugbayurdakadim@hotmail.com]