GENERALIZED ALMOST STARLIKENESS ASSOCIATED WITH EXTENSION OPERATORS FOR BIHOLOMORPHIC MAPPINGS

LUMINIŢA-IOANA COTÎRLĂ

Abstract. In this paper we consider an operator introduced by Pfaltzgraff and Suffridge which provides a way of extending a locally biholomorphic mapping $f \in H(B^n)$ to a locally biholomorphic mapping $F \in H(B^{n+1})$. Using the Loewner chains we prove that if f is an generalized almost starlike mapping on B^n then F is also generalized almost starlike mapping on B^{n+1} .

MSC 2010. 34C45,32H02.

Key words. Loewner chain, Roper-Suffridge extension operator, starlikeness, generalized almost starlikeness.

REFERENCES

- BĂLĂEȚI, C.M. and NECHITA, V.O., Loewner chains and almost starlike mappings of complex order λ, Carpathian J. Math., 26 (2010), 146–157.
- [2] COTÎRLĂ, L.I., Loewner chains and generalized almost starlike mappings, Mathematica, 54 (77), to appear.
- [3] FENG, S.X., Some classes of holomorphic mappings in several complex variables, University of Science and Technology of China, Doctoral thesis, 2004.
- [4] GRAHAM, I., KOHR, G., Geometric function theory in one and higher dimension, Marcel Dekker, New York, 2003.
- [5] GRAHAM, I., KOHR, G. and KOHR, M., Loewner chains and the Roper-Suffridge extension operator, J. Math. Anal. Appl., 247 (2000), 448–465.
- [6] GRAHAM, I., KOHR, G., KOHR, M., Loewner chains and parametric representation in several complex variables, J. Math. Anal. Appl., 281 (2003), 425–438.
- [7] GRAHAM, I., KOHR, G. and PFALTZGRAFF, J.A., Parametric representation and linear functionals associated with extension operators for biholomorphic mappings, Rev. Roumaine Math. Pure Appl., 52 (2007), 47–68.
- [8] GRAHAM, I., HAMADA, H. and KOHR, G., Parametric representation of univalent mappings in several complex variables, Canad. J.Math., 54 (2002), 324–351.
- [9] HAMADA,H. and KOHR, G., Subordination chains and the growth theorem of spirallike mappings, Mathematica, **42** (65) (2000), 153–161.
- [10] KOHR, G., On some sufficient conditions of almost starlikeness of order 1/2 in Cⁿ, Stud. Univ. Babes-Bolyai Math., 41 (1996), 51–55.
- [11] PFALTZGRAFF, J.A. and SUFFRIDGE, T.J., An extension theorem and linear invariant families generated by starlike maps, Ann. Univ. Mariae Curie-Sklodowska, Sect. A, 53 (1999), 193–207.
- [12] POREDA, T., On the univalent subordination chain of holomorphic mappings in Banach spaces, Comment. Math., 28 (1989), 295–304.
- [13] POREDA, T., On the univalent holomorphic maps of the unit polydisc of Cⁿ which have the parametric representation, II-necessary and sufficient conditions, Ann. Univ. Mariae Curie-Sklodowska, Sect. A, 41 (1987), 114–121.
- [14] ROPER, K. and SUFFRIDGE, T.J., Convexity properties of holomorphic mappings in \mathbb{C}^n , Trans. Amer. Math. Soc., **351** (1999), 1803–1833.

[15] RUDIN, W., Function Theory in the Unit Ball of Cⁿ, Springer-Verlag, New-York, 1980.
[16] XU, Q.-H. and LIU, T.S., Loewner chains and a subclass of biholomorphic mappings, J. Math. Anal. Appl., **334** (2007), 1096–1105.

Received January 15, 2010 Accepted May 20, 2010 "Babeş-Bolyai" University Faculty of Mathematics and Computer Science Str. Mihail Kogălniceanu Nr. 1 400084 Cluj-Napoca, Romania E-mail: uluminita@math.ubbcluj.ro E-mail: luminita.cotirla@yahoo.com