

ESTIMATES WITH OPTIMAL CONSTANTS USING PEETRE'S K -FUNCTIONALS OF ORDER 2

Maria Talpău Dimitriu

*Transilvania University, Department of Mathematics, Eroilor 29, 500
036, Braşov, Romania*

[marias_td@yahoo.com, mdimitriu@unitbv.ro]

2000 Mathematics Subject Classification. 41A36, 41A25.

Keywords and phrases. Positive linear operators, Peetre's K -functional, degree of approximation.

We present estimates of the degree of approximation by positive linear operators with the K -functionals K_2^s , $1 \leq s \leq \infty$.

REFERENCES

- [1] J. Bergh, J. Löfström, *Interpolation Spaces*, Springer-Verlag, 1976.
- [2] H. H. Gonska, *On approximation in spaces of continuous functions*, Bull. Austral. Math. Soc. 28 (1983), pp. 411-432.
- [3] H. H. Gonska, *Quantitative Korovkin type theorems on simultaneous approximation*, Math. Z. **186** (1984), pp. 419-433.
- [4] R. Păltănea, *Optimal estimates with moduli of continuity*, Result. Math. **32** (1997), pp. 318-331.
- [5] R. Păltănea, *Approximation theory using positive linear operators*, Birkhäuser, 2004.
- [6] J. Peetre, *On the connection between the theory of interpolation spaces and approximation theory*, Proc. Conf. Const. Theory of Functions, Budapest, Eds. G. Alexits and S. B. Stechkin, Akadémiai Kiadó, Budapest, 1969, pp. 351-363.
- [7] D. D. Stancu, *Quadrature formulas constructed by using certain linear positive operators*, Numerische Integration, 1982, pp. 241-251.