THE CONSTRUCTION OF COMPLETED
SKEW GROUP RINGS. II

IOAN FECHETE

Abstract. Given a profinite group G acting on a profinite ring R via $\sigma : G \to \mathrm{Aut}\,R$ (with finite image) we will construct the completition of the skew group ring $R * G$ as a inverse limit of skew group rings of type $R * (G/N)$, where N is a open invariant subgroup of G contained in $\ker \sigma$.

MSC 2000. 16W80.

Key words. Skew group ring, compact ring, profinite group (ring), completion of a topological ring.

REFERENCES

[1] FECHETE, I., *The construction of completed skew group rings*, Proc. Conf. Appl. Ind. Math., Oradea 2003, 100–103.

[2] FECHETE, I. and URSUL, M.I., *Completed commutative group rings without zero divisors*, Bul. Acad. Ştiinte Rep. Mold., Mat., **1(38)** (2002), 53–60.

[3] KOCH, H., *Galoissche Theorie der p-Erweiterungen*, Math. Monogr. **10**, VEB, Berlin, 1970.

[4] PASSMAN, D.S., *Group Rings, Crossed Products and Galois Theory*, Regional Conference Series in Mathematics, **64**, A.M.S., Providence, RI, 1986.

[5] URSUL, M.I., *Compact rings satisfying compactness conditions*, Kluwer, Doodrecht, 2002.

University of Oradea
Department of Mathematics
str. Armatei Romane, 5
410087, Oradea, ROMANIA
E-mail: `ifechete@uoradea.ro`