MATHEMATICA, Tome 47 (70), N° 1, 2005, pp. 61–72

OPENNESS RESULTS FOR MULTIFUNCTIONS VIA AN ABSTRACT SUBDIFFERENTIAL

MARIUS DUREA

Abstract. In this paper we establish some openness-type results for set-valued maps in Banach spaces, using a coderivarive which is defined via abstract subdifferentials. Mainly, results of openness at a linear rate are obtained. Then we apply these results to derive a necessary optimality condition for a nonlinear optimization problem, without any convexity assumptions.

MSC 2000. 46T99, 54C60, 90C26.

Key words. Abstract subdifferential, multifunction, openness at a linear rate, coderivative, optimization formula.

REFERENCES

- [1] AUBIN, J.-P. and FRANKOWSKA, H., *Set-valued Analysis*, Birkhäuser, Basel, Berlin, 1990.
- [2] AZÉ, D., CHOU, C.C. and PENOT, J.-P., Subtraction theorems and approximate openness for multifunctions: topological and infinitesimal viewpoints, J. Math. Anal. Appl., 221 (1998), 33–58.
- [3] BORWEIN, J.M. and ZHU, Q.J., Viscosity solutions and viscosity subderivatives in smooth Banach spaces with applications to metric regularity, SIAM J. Control Optim., 34 (1996), 1568–1591.
- [4] CORLEY, H.W., Optimality conditions for maximization of set-valued functions, J. Optim. Theory Appl., 58 (1988), 1–10.
- [5] IOFFE, A.D., Approximate subdifferentials and applications. I: The finite dimensional theory, Trans. Amer. Math. Soc., 281 (1984), 389–416.
- [6] IOFFE, A.D., Metric regularity and subdifferential calculus, Russian Math. Surveys, 55 (2000), 501–558.
- [7] IOFFE, A.D. and PENOT, J.-P., Subdifferentials of performance functions and calculus of coderivatives of set-valued mappings, Serdica Math. J., 22 (1996), 359–384.
- [8] JOURANI, A., Regularity and strong sufficient optimality conditions in differentiable optimization problems, Numer. Funct. Anal. and Optim., 14 (1993), 69–87.
- [9] KLEIN, E. and THOMPSON, A.C., *Theory of Correspondences*, Wiley-Interscience Publication, 1984.
- [10] LI, Z., A Theorem of the alternative and its application to the optimization of set-valued maps, J. Optim. Theory Appl., 100 (1999), 365–375.
- [11] MORDUKHOVICH, B., Complete characterization of openness, metric regularity, and lipschitzian behavior of multifunctions, Trans. Amer. Math. Soc., 340 (1993), 1–35.
- [12] PENOT, J.-P., Cooperative behavior of functions, relations and sets, Math. Methods Oper. Res., 48 (1998), 229–246.

Received September 17, 2003

Universitatea "Al. I. Cuza" Iași Facultatea de Matematică Bd. Carol I, Nr. 11 700506 Iași, România E-mail: durea@uaic.ro