

SUFFICIENT CONDITIONS FOR STARLIKENESS
AND CONVEXITY
OF ANALYTIC FUNCTIONS WITH NEGATIVE COEFFICIENTS

G.S. SĂLĂGEAN

Abstract. Several interesting implications concerning analytic functions with negative coefficients are determined. In particular cases sufficient conditions for starlikeness, strongly starlikeness and convexity are obtained.

MSC 2000. 30C45.

Key words. Starlike, convex, strongly starlike, Libera operator.

REFERENCES

- [1] S. S. MILLER and P. T. MOCANU, *Libera transform of functions with bounded turning*, J. Math. Anal. Appl., **276** (2002), 90–97.
- [2] P. T. MOCANU *On starlikeness of Libera transform*, Mathematica (Cluj), **28 (51)** (1986), 153–155.
- [3] P. T. MOCANU, *New extensions of a theorem of R. Singh and S. Singh*, Mathematica (Cluj), **37 (60)** (1995), 171–182.
- [4] R. SINGH AND S. SINGH, *Starlikeness and convexity of certain integral*, Ann. Univ. Marie Curie-Sklodowska, Sect. A, **35** (1981), 145–148.
- [5] H. SILVERMAN, *Univalent functions with negative coefficients*, Proc. Amer. Math. Soc. **51** (1975), 109–116.
- [6] H. SILVERMAN, *A survey with open problems on univalent functions whose coefficients are negative*, Rocky Mountain J. Math., **21** (1991), 3, 1099–1125.
- [7] G. S. SĂLĂGEAN, *Subclasses of univalent functions*, Complex Analysis, Fifth Romanian-Finnish Sem., Lect. Notes in Math., **1013**, Springer-Verlag, 1983, 362–372.
- [8] G. S. SĂLĂGEAN, *Classes of univalent functions with two fixed points*, “Babeş-Bolyai” Univ., Res. Sem., 6/1994, 181–184.
- [9] G. S. SĂLĂGEAN, *On univalent functions with negative coefficients*, “Babeş-Bolyai” Univ., Res. Sem., 7/1991, 47–54.

Received December 15, 2004

*Faculty of Mathematics and Computer Science
“Babeş-Bolyai” University
1 M. Kogălniceanu Str.
400084 Cluj-Napoca, Romania
E-mail: salagean@math.ubbcluj.ro*