SOME INTEGRAL OPERATORS DEFINED ON *p*-VALENT FUNCTIONS BY USING HYPERGEOMETRIC FUNCTIONS

A. TEHRANCHI and S. R. KULKARNI

Abstract. In the present paper we introduce some integral operators and verify the effect of these operators on *p*-valent functions and find radii of starlikeness and convexity for these operators, finally we introduce the concept of neighborhood.

MSC 2000. 30C45, 30C50.

Key words. Uniformly convex functions, fractional derivative, starlike and convex p-valent, hypergeometric, neighborhood.

REFERENCES

- ALTINTAS, O. On a subclass of certain starlike functions with negative coefficient, Math. Japonica, 36 (1991), 489–495.
- [2] ALTINTAS, O. and OWA, S., Neighborhood of certain analytic function with negative coefficients, Int. J. Math. Sci., 19 (1996), 797–800.
- [3] AQLAN, E., Some problems connected with geometric function theory, Ph.D. Thesis, University of Pune, India, 2004.
- [4] ANDREWS, G.E., ASKEY, R. and ROY, R. Special functions, Cambridge University Press, Cambridge, UK, (1991).
- [5] BHARATI, R., PARVATHAM, R. and SWAMINATHAN, A., On subclasses of uniformly convex functions and corresponding class of starlike functions, Tamkang, J. Math., 28 (1997), 17–32.
- [6] KANAS, S. and WIŚNIOWSKA, A., Conic regions and k-uniform convexity, J. Comp. Appl. Math., 105 (1999), 327–336.
- [7] KIM, Y.C. and RØNNING, F. Integral transforms of certain subclasses of analytic functions, Jour. Math. Anal. Appl., 258 (2001), 466–489.
- [8] KOMATO, Y. On analytic prolongation of a family of operators, Mathematica (Cluj), 55 (1990), 141–145.
- [9] RUCHEWEYH, S. Neighborhoods of univalent functions, Proc. Amer. Mth. Soc., 81 (1981), 521–527.
- [10] SILVERMAN, H. Univalent functions with negative coefficients, Proc. Amer. Math. Soc., 51 (1975), 109–116.
- [11] SRIVASTAVA, H.M. and OWA, S. An application of the fractional derivative, Math. Japonica, **29** (1984), 383–389.
- [12] SRIVASTAVA, H. M., OWA, S. Univalent functions fractional calculus and their applications. Halsted Press (Ellis Horwood Limited Chichester), John Wiley and Sons, New York, Chichester Brisbane and Toronto (1989).
- [13] TEHRANCHI, A. and KULKARNI, S. R., Subclass of α -uniformly starlike and convex pvalent functions in terms of fractional derivative of f(z) of order δ (I) (submitted for publication).

Received March 5, 2005

Department of Mathematics, Fergusson College, University of Pune, Pune - 411004, India E-mail: kulkarni_ferg@yahoo.com E-mail: Tehranchiab@yahoo.co.uk