THE REDUCTION OF A k-SYMPLECTIC MANIFOLD

ADARA M. BLAGA

Abstract. The aim of this paper is to construct an analogue of Marsden-Weinstein reduction for k-symplectic manifolds.

MSC 2000. 53C15.

Key words. Momentum map, symplectic manifold, Marsden-Weinstein reduction.

REFERENCES

[1] Abraham, R. and Marsden, J., *Foundations of mechanics*, Benjamin, New York, (1967).

[2] Awane, A., *k-symplectic structures*, J. Math. Phys., **33** (1992), 4046–4052.

[3] Cantrijn, A. and De Leon, M., *On the geometry of multisymplectic manifolds*, preprint IMAFF (1996).

[4] Cantrijn, A. and De Leon, M., *Hamiltonian structures on multisymplectic manifolds*, Proceedings of the Workshop on Geometry and Phisics, Italy (1996).

[5] Cendra, H., Marsden, J. and Raţiu, T., *Reduction, symmetry and phases in mechanics*, An. Math. Soc. (1990).

[6] Günther, Ch., *The polysymplectic Hamiltonian formalism in field theory and calculus of variations*, J. Diff. Geom., **25** (1987), 23–53.

[7] Martin, G., *A Darboux theorem for multisymplectic manifolds*, Lett. Math. Phys., **16** (1988), 133–138.

[8] Munteanu, F., Rey, A. and Salgado, M., *The Günther's formalism in classical field theory: momentum map and reduction*, J. Math. Phys., **45** (2004), 1730–1751.

[9] Norris, L.K., *Symplectic geometry on T^*M derived from k-symplectic geometry on LM*, J. Geom. Phys., **13** (1994), 51–78.

[10] Puta, M., *Some remarks on k-symplectic manifolds*, Tensor, **47** (1998), 109–115.

West University from Timişoara
Departament of Mathematics and Computer Sciences
Bdl. V. Pârvan nr.4, 300223 Timişoara, România
E-mail: `adara@math.uvt.ro`