$Q_{K,\omega,\log}(p,q)$ -TYPE SPACES OF ANALYTIC AND MEROMORPHIC FUNCTIONS

A. EL-SAYED AHMED and ALAA KAMAL

Abstract. In this paper, we define the space $Q_{K,\omega,\log}(p,q)$ of analytic functions on the unit disk. We obtain some characterizations for the space $Q_{K,\omega,\log}(p,q)$ by the help of the nondecreasing function K and the reasonable function ω . Moreover, the meromorphic $Q_{K,\omega,\log}^{\#}(p,q)$ space is also considered and studied.

MSC 2010. 30D45, 46E15.

Key words. Logarithmic Bloch functions, $Q_{K,\omega,\log}(p,q)$ spaces.

REFERENCES

- [1] EL-SAYED AHMED, A. and BAHKIT, M.A., Hadamard products and \mathcal{N}_K space, Math. Comput. Modelling, **51** (2010), 33–43.
- [2] EL-SAYED AHMED, A. and BAHKIT, M.A., Sequences of some meromorphic function spaces, Bull. Belg. Math. Soc. Simon Stevin, 16 (2009), 395–408.
- [3] AULASKARI, R. and LAPPAN, P., Criteria for an analytic function to be Bloch and a harmonic or meromorphic function to be normal. In Complex Analysis and its Applications (Y. Chung-Chun et al., Eds.), Pitman Research Notes in Mathematics, 305, Longman, 1994, pp. 136–146.
- [4] DYAKONOV, K.M., Weighted Bloch spaces, H^p, and BMOA, J. Lond. Math. Soc. (2), 65 (2002), 411–417.
- [5] ESSÉN, M. and WULAN, H., On analytic and meromorphic functions and spaces of Q_K type, Illinois J. Math., **46** (2002), 1233–1258.
- [6] ESSÉN, M. WULAN, H. and XIAO, J., Several Function-theoretic characterizations of Möbius invariant Q_K spaces, J. Funct. Anal., 230 (2006), 78–115.
- [7] LAPPAN, P., A non-normal locally uniformly univalent function, Bull. London Math. Soc., 5 (1973), 491–495.
- [8] RASHWAN, R.A., EL-SAYED AHMED, A. and KAMAL, A., Some characterizations of weighted holomorphic Bloch space, Eur. J. Pure Appl. Math., 2 (2009), 250–267.
- [9] RASHWAN, R.A., EL-SAYED AHMED, A. and KAMAL, A., Integral characterizations of weighted Bloch space and $Q_{K,\omega}(p,q)$ spaces, Mathematica, **51** (74) (2009), 63–76.
- [10] RÄTTYÄ, J., On some Complex function spaces and classes, Ann. Acad. Sci. Fenn. Math. Diss., 124 (2001), 1–73.
- [11] STROETHOFF, K., Besov-type characterisations for the Bloch space, Bull. Austral. Math. Soc., 39 (1989), 405–420.
- [12] WULAN, H., On some classes of meromorphic functions, Ann. Acad. Sci. Fenn. Math. Diss., 116 (1998), 1–57.
- [13] WULAN, H. and WU, P., Characterizations of Q_T spaces, J. Math. Anal. Appl., 254 (2001), 484–497.
- [14] WULAN, H. and ZHU, K., Q_K type spaces of analytic functions, J. Funct. Spaces Appl., 4 (2006), 73–84.
- [15] YONEDA, R., The composition operators on weighted Bloch space, Arch. Math. (Basel), 78 (2002), 310–317.

Received December 30, 2010 Accepted April 2, 2012 Sohag University Faculty of Science Mathematics Department 82524 Sohag, Egypt

Taif University Faculty of Science Mathematics Department El-Taif 5700, El-Hawiyah, Saudi Arabia E-mail: ahsayed80@hotmail.com

Majmaah University Faculty of Science Department of Mathematics Saudi Arabia E-mail: alaa_mohamed1@yahoo.com