MATHEMATICA, 62 (85), N $^{\circ}$ 1, 2020, pp. 33–38

RECTANGULAR TRANSFORMATIONS IN LATIN SQUARES

IVAN I. DERIYENKO

Abstract. To get another one from a given latin square, we have to change at least 4 entries. We show how to find these entries and how to change them. MSC 2010. 05B15, 20N05.

Key words. Autotopism, latin square, quasigroup.

REFERENCES

- V.D. Belousov, The group associated with a quasigroup (in Russian), Mat. Issled., 4 (1969), 21–39.
- [2] I.I. Deriyenko, Necessary conditions of the isotopy of finite quasigroups (in Russian), Mat. Issled., 120 (1991), 51–63.
- [3] I.I. Deriyenko, *Indicators of quasigroups*, Quasigroups Related Systems, 19 (2011), 223– 226.
- [4] I.I. Deriyenko and W.A. Dudek, On prolongations of quasigroups, Quasigroups Related Systems, 16 (2008), 187–198.
- [5] I.I. Deriyenko and W.A. Dudek, *D-loops*, Quasigroups Related Systems, 20 (2012), 183–196.
- [6] D. Donovan, S. Oates-Williams and Ch. E. Praeger, On the distance between distinct group latin squares, J. Combin. Des., 5 (1997), 235–248.
- [7] A. Drápal, Hamming distances of groups and quasigroups, Discrete Math., 235 (2001), 189–197.
- [8] W.A. Dudek, Parastrophes of quasigroups, Quasigroups Related Systems, 23 (2015), 221–230.
- [9] S.A. Frisch, On the minimal distance between group tables, Acta Sci. Math. (Szeged), 63 (1997), 341–351.
- [10] D. Keedwell and J. Dénes, Latin squares and their applications, North-Holland, 2015.
- [11] V. Shcherbacov, Elements of quasigroup theory and applications, CRC Press, 2017.

Received January 23, 2019Kremenchuk State Polytechnic UniversityAccepted March 14, 2019Department of Higher Mathematics and Informatics
39600 Kremenchuk, Ukraine
E-mail: ivan.deriyenko@gmail.com

DOI: 10.24193/mathcluj.2020.1.04