

# Lecture 6

# Chapter 4. Numerical Characteristics of Random Variables

- the distribution of a random variable or a random vector, the full collection of related probabilities, contains the entire information about its behavior;
- this detailed information can be summarized in a few **vital numerical characteristics** describing the *average value, the most likely value of a random variable, its spread, variability, etc*;
- these are **numbers** that will provide some information about a random variable or about the relationship between random variables.

# 1. Expectation

## Definition 1.1.

- (i) If  $X \left( \begin{array}{c} x_i \\ p_i \end{array} \right)_{i \in I}$  is a discrete random variable, then the **expectation (expected value, mean value)** of  $X$  is the real number

$$E(X) = \sum_{i \in I} x_i P(X = x_i) = \sum_{i \in I} x_i p_i, \quad (1.1)$$

if it exists (i.e., the series is absolutely convergent).

- (ii) If  $X$  is a continuous random variable with density function  $f : \mathbb{R} \rightarrow \mathbb{R}$ , then its **expectation (expected value, mean value)** is the real number

$$E(X) = \int_{\mathbb{R}} xf(x)dx, \quad (1.2)$$

if it exists (i.e., the integral is absolutely convergent).

**Remark 1.2.**

1. The expected value can be thought of as a “**long term**” **average value**, a number that we *expect* the values of a random variable to stabilize on.
2. If  $h : \mathbb{R} \rightarrow \mathbb{R}$  is a measurable function, then

$$E(h(X)) = \sum_{i \in I} h(x_i) p_i, \quad (1.3)$$

if  $X$  is discrete and

$$E(h(X)) = \int_{\mathbb{R}} h(x) f(x) dx, \quad (1.4)$$

if  $X$  is continuous.

It can also be interpreted as a **point of equilibrium**, a **center of gravity**.

In the discrete case, if we imagine the probabilities  $p_i$  to be **weights** distributed in the points  $x_i$ , then  $E(X)$  would be the point that holds the whole thing in *equilibrium*. In fact, notice that the computational formula (1.1) is *actually* a weighted mean.

Consider a random variable with pdf

$$X \left( \begin{array}{cc} 0 & 1 \\ 0.5 & 0.5 \end{array} \right).$$

Observing this variable many times, we shall see  $X = 0$  about 50% of times and  $X = 1$  about 50% of times. The average value of  $X$  will then be close to 0.5, so it is reasonable to have  $E(X) = 0.5$ , which is what we get by (1.1).

Now, suppose that  $P(X = 0) = 0.75$  and  $P(X = 1) = 0.25$ , i.e its pdf is now


$$X \begin{pmatrix} 0 & 1 \\ 0.75 & 0.25 \end{pmatrix}.$$

Then, in a long run,  $X$  is equal to 1 only 1/4 of times, otherwise it equals 0. Therefore, in this case,  $E(X) = 0.25$ .

The same interpretation would go for the continuous case, only there the “weight” would be continuously distributed, according to the density function  $f$ .

The expected value as a center of gravity is illustrated in Figure 1.

$$(a) \mathbf{E}(X) = 0.5$$



$$(b) \mathbf{E}(X) = 0.25$$




Figure 1: Expectation as a center of gravity

**Example 1.3.**

Let us start with a simple, intuitive example. Let  $X$  be the random variable that denotes the number shown when a die is rolled. What would be the “expected average value” of  $X$ , if the die was rolled over and over?

**Solution.**

Since any of the 6 numbers is **equally probable** to show on the die, we would expect that, in the long run, we would roll **as many 1’s as 6’s**. These would average out at

$$\frac{1 + 6}{2} = \frac{7}{2}.$$

Also, we would expect to roll **the same number of 2’s as 5’s**, which would also average at

$$\frac{2 + 5}{2} = \frac{7}{2}.$$

Finally, about **the same number of 3’s and 4’s** would be expected to show and their average is again,  $\frac{7}{2}$ . So, the “long term average” should be, intuitively,  $\frac{7}{2}$ .



On the other hand, we know that  $X$  has a **Discrete Uniform**  $U(6)$  distribution, with pdf

$$X \left( \begin{array}{cccccc} 1 & 2 & 3 & 4 & 5 & 6 \\ \frac{1}{6} & \frac{1}{6} & \frac{1}{6} & \frac{1}{6} & \frac{1}{6} & \frac{1}{6} \end{array} \right).$$

Then, by (1.1),

$$\begin{aligned} E(X) &= \sum_{i \in I} x_i p_i \\ &= \sum_{i=1}^6 i \cdot \frac{1}{6} = \frac{1}{6} \sum_{i=1}^6 i \\ &= \frac{1}{6} \cdot \frac{6 \cdot 7}{2} = \frac{7}{2}, \end{aligned}$$

the value we obtained intuitively.



### Example 1.4.

Consider now a (continuous) Uniform variable  $X \in U(a, b)$ . That means  $X$  can take *any* value in the interval  $[a, b]$ , **equally probable** (recall Problem 3 in Seminar 2, about a spyware breaking passwords). What would be a long-run “expected average value”?

### Solution.

In the long run, the variable is just as likely to take values at the beginning of the interval, as it is to take the ones towards the end of  $[a, b]$ . So they would average out at the value right in the middle, i.e. the **midpoint** of the interval,

$$\frac{a + b}{2}.$$

Indeed, since the pdf of  $X$  is

$$f(x) = \frac{1}{b-a}, x \in [a, b]$$

(and 0 everywhere else), by (1.2), its expected value is

$$\begin{aligned} E(X) &= \int_{\mathbb{R}} xf(x)dx = \int_a^b x \frac{1}{b-a} dx = \frac{1}{b-a} \int_a^b x dx \\ &= \frac{1}{b-a} \cdot \frac{1}{2} x^2 \Big|_a^b = \frac{1}{b-a} \cdot \frac{b^2 - a^2}{2} = \frac{a+b}{2}. \end{aligned}$$



**Example 1.5.**

The expected value of a  $Bern(p)$ ,  $p \in (0, 1)$  variable with pdf

$$X \begin{pmatrix} 0 & 1 \\ 1-p & p \end{pmatrix}$$

is

$$E(X) = 0 \cdot (1-p) + 1 \cdot p = p. \quad (1.5)$$

**Theorem 1.6.**

If  $X$  and  $Y$  are either both discrete or both continuous random variables, then the following properties hold:

- a)  $E(aX + b) = aE(X) + b$ , for all  $a, b \in \mathbb{R}$ .
- b)  $E(X + Y) = E(X) + E(Y)$ .
- c) If  $X$  and  $Y$  are independent, then  $E(X \cdot Y) = E(X)E(Y)$ .
- d) If  $X \leq Y$ , i.e.  $X(e) \leq Y(e)$ , for all  $e \in S$ , then  $E(X) \leq E(Y)$ .

## Proof.

We give a selected proof, **only for the discrete case.**

a) If  $X$  is discrete, with pdf

$$X \left( \begin{array}{c} x_i \\ p_i \end{array} \right)_{i \in I},$$

then  $Y = aX + b$  is also discrete and has pdf

$$Y \left( \begin{array}{c} ax_i + b \\ p_i \end{array} \right)_{i \in I}.$$

So, its expectation is

$$E(aX + b) = \sum_{i \in I} (ax_i + b)p_i = a \sum_{i \in I} x_i p_i + b \sum_{i \in I} p_i = aE(X) + b.$$

Proof.

b) For  $X$  and  $Y$  both discrete, recall that their **sum** has pdf

$$X + Y \left( \begin{array}{c} x_i + y_j \\ p_{ij} \end{array} \right)_{(i,j) \in I \times J}, \quad p_{ij} = P(X = x_i, Y = y_j)$$

and that

$$\sum_{j \in J} p_{ij} = p_i, \quad \sum_{i \in I} p_{ij} = q_j,$$

where  $p_i = P(X = x_i)$ ,  $i \in I$  and  $q_j = P(Y = y_j)$ ,  $j \in J$ .

Proof.

Then

$$\begin{aligned}
 E(X + Y) &= \sum_{(i,j) \in I \times J} (x_i + y_j) p_{ij} = \sum_{i \in I} \sum_{j \in J} (x_i + y_j) p_{ij} \\
 &= \sum_{i \in I} \sum_{j \in J} x_i p_{ij} + \sum_{j \in J} \sum_{i \in I} y_j p_{ij} \\
 &= \sum_{i \in I} x_i \underbrace{\sum_{j \in J} p_{ij}}_{p_i} + \sum_{j \in J} y_j \underbrace{\sum_{i \in I} p_{ij}}_{q_j} \\
 &= \sum_{i \in I} x_i p_i + \sum_{j \in J} y_j q_j \\
 &= E(X) + E(Y).
 \end{aligned}$$



Proof.

c) For  $X$  and  $Y$  discrete and independent, we have

$$\begin{aligned}
 E(XY) &= \sum_{i \in I} \sum_{j \in J} x_i y_j p_{ij} \stackrel{\text{ind}}{=} \sum_{i \in I} \sum_{j \in J} x_i y_j p_i q_j \\
 &= \sum_{i \in I} x_i \underbrace{\left( \sum_{j \in J} y_j q_j \right)}_{E(Y)} p_i \\
 &= E(Y) \cdot \sum_{i \in I} x_i p_i \\
 &= E(X) \cdot E(Y).
 \end{aligned}$$

Proof.

**d)** We show that if  $Z \geq 0$ , then  $E(Z) \geq 0$ .

Then by a) and b) applied to  $Z = Y - X$ , the property follows.

If  $Z$  is discrete,  $Z \geq 0$  means **its values**  $z_i \geq 0$ ,  $\forall i \in I$  and then

$$E(Z) = \sum_{i \in I} z_i P(Z = z_i) \geq 0.$$



**Remark 1.7.**

1. Property b) in Theorem 1.6 can be generalized to

$$E\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n E(X_i).$$

2. Property c) in Theorem 1.6 can also be generalized: If  $X_1, \dots, X_n$  are **independent**, then

$$E\left(\prod_{i=1}^n X_i\right) = \prod_{i=1}^n E(X_i).$$

3. An immediate consequence of Theorem 1.6a) is the fact that

$$E(X - E(X)) = 0.$$

**Example 1.8.**

Let us find the expectation of a Binomial variable  $X \in B(n, p)$ ,  $n \in \mathbb{N}$ ,  $p \in (0, 1)$ .

**Solution.** Recall (Remark 4.8, Lecture 4) that a Binomial variable  $X \in B(n, p)$  is the sum of  $n$  independent  $X_i \in \text{Bern}(p)$  random variables. All variables  $X_i$  have the same expected value  $E(X_i) = p$ , since they have the same distribution.

Then, by the previous theorem,

$$E(X) = E\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n E(X_i) = \sum_{i=1}^n p = np.$$

**Remark 1.9.**

For a Normal variable  $X \in N(\mu, \sigma)$ , the expected value is  $E(X) = \mu$ .

## 2. Variance and Standard Deviation

Expectation shows where the average value of a random variable is located, or where the variable is expected to be, plus or minus **some error**. How **large** could this “error” be, and how much can a variable vary around its expectation? The answer to these questions can give important information about a random variable.

Knowledge of the mean value of a random variable is important, but that knowledge *alone* can be misleading. Suppose two patients in a hospital,  $X$  and  $Y$ , have their pulse (number of heartbeats per minute) checked every day. Over the course of time, they each have a mean pulse of 75, which is considered healthy. But, for patient  $X$  the pulse ranges between 70 and 80, while for patient  $Y$ , it oscillates between 40 and 110. Obviously, the second patient might have some serious health problems, which the *expected value alone* would not show.

So, next, we define some measures of **variability**.

## Definition 2.1.

Let  $X$  be a random variable. The **variance (dispersion)** of  $X$  is the number

$$V(X) = E\left[\left(X - E(X)\right)^2\right], \quad (2.1)$$

if it exists. The value  $\sigma(X) = \text{Std}(X) = \sqrt{V(X)}$  is called the **standard deviation** of  $X$ .

Variance (and standard deviation) measure the amount of **variability** (spread) in the values that a random variable takes, with **large values** indicating a **wide spread** of values and **small values** meaning **more closely knit** values.

The standard deviation brings the numbers to the same “level” (e.g., measurement units), while the variance gives the squares of those numbers.

**Theorem 2.2.**

Let  $X$  and  $Y$  be random variables. Then the following properties hold:

- a)  $V(X) = E(X^2) - (E(X))^2$ .
- b)  $V(aX + b) = a^2V(X)$ , for all  $a, b \in \mathbb{R}$ .
- c) If  $X$  and  $Y$  are independent, then

$$V(X + Y) = V(X) + V(Y).$$

- d) If  $X$  and  $Y$  are independent, then

$$V(X \cdot Y) = E(X^2)E(Y^2) - (E(X))^2(E(Y))^2.$$

## Proof.

We give a selected proof.

a) By properties of expectation in Theorem 1.6, we have

$$\begin{aligned}V(X) &= E\left[X^2 - 2E(X)X + (E(X))^2\right] \\&= E(X^2) - 2E(X)E(X) + (E(X))^2 \\&= E(X^2) - (E(X))^2.\end{aligned}$$

b)

$$\begin{aligned}V(aX + b) &= E\left[(aX + b - E(aX + b))^2\right] \\&= E\left[(aX + b - aE(X) - b)^2\right] \\&= a^2E\left[(X - E(X))^2\right] = a^2V(X).\end{aligned}$$



Proof.

c) If  $X, Y$  are **independent**, then **so are**  $X - E(X), Y - E(Y)$ , thus,

$$\begin{aligned}
 V(X + Y) &= E\left[(X + Y - E(X + Y))^2\right] \\
 &= E\left[((X - E(X)) + (Y - E(Y)))^2\right] \\
 &= E\left[(X - E(X))^2\right] + 2E\left[(X - E(X))(Y - E(Y))\right] \\
 &\quad + E\left[(Y - E(Y))^2\right] \\
 &\stackrel{\text{ind}}{=} V(X) + 2E(X - E(X)) \cdot E(Y - E(Y)) + V(Y) \\
 &= V(X) + V(Y),
 \end{aligned}$$

since  $E(X - E(X)) = 0$ .



**Remark 2.3.**

1. Part a) of Theorem 2.2 provides a **more practical computational formula** for the variance than the definition.

Thus, if  $X \left( \begin{array}{c} x_i \\ p_i \end{array} \right)_{i \in I}$  is discrete, then

$$V(X) = \sum_{i \in I} x_i^2 p_i - \left( \sum_{i \in I} x_i p_i \right)^2$$

and if  $X$  is continuous with density function  $f$ , then

$$V(X) = \int_{\mathbb{R}} x^2 f(x) dx - \left( \int_{\mathbb{R}} x f(x) dx \right)^2.$$

**Remark 2.3.**

2. A direct consequence of Theorem 2.2a) (since  $V(X) \geq 0$ ) is the following inequality:

$$|E(X)| \leq \sqrt{E(X^2)},$$

which will be discussed later on in this chapter.

3. If  $X = b$  is a constant random variable (i.e. it only takes that one value with probability 1), then by Theorem 2.2a),  $V(X) = 0$ , which is to be expected (the variable  $X$  **does not vary at all**).

**Remark 2.3.**

4. Part c) of Theorem 2.2 can be generalized to **any number** of random variables: If  $X_1, \dots, X_n$  are independent, then

$$V\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n V(X_i).$$

5. A consequence of parts b) and c) of Theorem 2.2 is the following property: If  $X$  and  $Y$  are independent, then

$$V(X + Y) = V(X) + V(Y) = V(X) + V(-Y) = V(X - Y).$$

**Example 2.4.**

Find the variance of a random variable  $X$  having

- a) a Bernoulli  $Bern(p)$  distribution;
- b) a Binomial  $B(n, p)$  distribution.

**Solution.**

a) We have

$$X \begin{pmatrix} 0 & 1 \\ 1-p & p \end{pmatrix}, \quad X^2 \begin{pmatrix} 0 & 1 \\ 1-p & p \end{pmatrix},$$

so both  $E(X) = E(X^2) = p$  and thus,

$$V(X) = p - p^2 = pq.$$

b) If  $X$  is Binomial, again we use the fact that it can be written as

$$X = \sum_{i=1}^n X_i,$$

where  $X_1, \dots, X_n$  are **independent and identically distributed** with a  $Bern(p)$  distribution. Then by part a),  $V(X_i) = pq$ , for each  $i = \overline{1, n}$  and by the previous remarks,

$$V(X) = V\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n V(X_i) = npq.$$

### Remark 2.5.

For a Normal variable  $X \in N(\mu, \sigma)$ , the variance is  $V(X) = \sigma^2$  and its standard deviation is  $\sigma(X) = \text{Std}(X) = \sigma$ . So, the parameters of a Normal variable  $X \in N(\mu, \sigma)$  are its **mean value** and its **standard deviation**.

### 3. Median

#### Definition 3.1.

The **median** of a random variable  $X$  with cdf  $F : \mathbb{R} \rightarrow \mathbb{R}$  is a real number  $M$  that is exceeded with probability no greater than 0.5 and is preceded with probability no greater than 0.5. That is,  $M$  is such that

$$P(X > M) \leq 1/2, \text{ i.e. } 1 - F(M) \leq 1/2,$$

$$P(X < M) \leq 1/2, \text{ i.e. } F(M - 0) \leq 1/2.$$

Comparing the mean  $E(X)$  and the median  $M$ , one can tell whether the distribution of  $X$  is

- right-skewed ( $M < E(X)$ ),
- left-skewed ( $M > E(X)$ ), or
- symmetric ( $M = E(X)$ ).

For *continuous* distributions, since  $P(X < M) = P(X \leq M) = F(M) = F(M - 0)$ , computing a population median reduces to solving one equation:

$$\begin{cases} P(X > M) = 1 - F(M) \leq 1/2 \\ P(X < M) = F(M) \leq 1/2 \end{cases} \Rightarrow F(M) = 1/2.$$

The Uniform distribution  $U(a, b)$  has cdf  $F(x) = \frac{x - a}{b - a}$ ,  $x \in [a, b]$ . Solving the equation  $F(M) = (M - a)/(b - a) = 1/2$ , we find the median

$$M = \frac{a + b}{2},$$

which is also the expected value  $E(X)$ . That should be no surprise, knowing that the Uniform distribution is **symmetric** (see Figure 2(a)).

For the Exponential distribution  $\text{Exp}(\lambda)$ , the cdf is  $F(x) = 1 - e^{-\lambda x}$ ,  $x > 0$ . Solving  $F(M) = 1 - e^{-\lambda M} = 1/2$ , we get

$$M = \frac{\ln 2}{\lambda} \approx \frac{0.6931}{\lambda} < \frac{1}{\lambda} = E(X),$$

since the Exponential distribution is **right-skewed** (see Figure 2(b)).



(a) Uniform

(b) Exponential






Figure 2: Median for Continuous Distributions

For *discrete* distributions, the equation  $F(x) = 0.5$  has either a **whole interval of roots** or **no roots** at all (see Figure 3).

(a) Binomial ( $n=5, p=0.5$ )  
many roots



(b) Binomial ( $n=5, p=0.4$ )  
no roots




Figure 3: Median for Discrete Distributions

In the first case, the Binomial distribution  $B(5, 0.5)$ , with  $p = 0.5$ , successes and failures are **equally likely**. Pick, for example,  $x = 2.2$  in the interval  $(2, 3)$ . Having **fewer** than 2.2 successes (i.e., at most 2) has the same chance as having **more** than 2.2 successes (i.e., at least 3 successes). Therefore,  $X < 2.2$  with the same probability as  $X > 2.2$ , which makes  $x = 2.2$  a central value, a **median**. The same is true for *any other* value in  $(2, 3)$ . In this case, for any number  $M \in (2, 3)$ , we have  $F(M) = F(M - 0) = 0.5$ , so any such number is a median.

In the other case, the Binomial distribution  $B(5, 0.4)$  with  $p = 0.4$ , we have  $F(1) = 0.2333$  and  $F(2) = 0.5443$ , so

$$\begin{aligned} F(x) &< 0.5 & \text{for } x < 2, \\ F(x) &> 0.5 & \text{for } x \geq 2, \end{aligned}$$

but there is **no value** of  $x$  with  $F(x) = 0.5$ . Then,  $M = 2$  is the **median**. Seeing a value on either side of  $M = 2$  has probability less than 0.5, which makes  $M = 2$  a center value. Here,  $F(M) > 0.5$  and  $F(M - 0) < 0.5$ .

## 4. Moments

The ideas of expected value and variance can be generalized.

### Definition 4.1.

Let  $X$  be a random variable and let  $k \in \mathbb{N}$ .

The **(initial) moment of order  $k$**  of  $X$  is (if it exists) the number

$$\nu_k = E(X^k). \quad (4.1)$$

The **absolute moment of order  $k$**  of  $X$  is (if it exists) the number

$$\underline{\nu}_k = E(|X|^k). \quad (4.2)$$

The **central (centered) moment of order  $k$**  of  $X$  is (if it exists) the number

$$\mu_k = E\left[(X - E(X))^k\right]. \quad (4.3)$$

**Remark 4.2.**

1. If  $X$  is a discrete random variable with pdf  $\left( \begin{matrix} x_i \\ p_i \end{matrix} \right)_{i \in I}$ , then for every  $k \in \mathbb{N}$ ,

$$\nu_k = \sum_{i \in I} x_i^k p_i, \quad \underline{\nu}_k = \sum_{i \in I} |x_i|^k p_i, \quad \mu_k = \sum_{i \in I} (x_i - E(X))^k p_i.$$

If  $X$  is a continuous random variable with pdf  $f$ , then for every  $k \in \mathbb{N}$ ,

$$\nu_k = \int_{\mathbb{R}} x^k f(x) dx, \quad \underline{\nu}_k = \int_{\mathbb{R}} |x|^k f(x) dx, \quad \mu_k = \int_{\mathbb{R}} (x - E(X))^k f(x) dx.$$

## Remark 4.2.

2. The **expectation** of a random variable  $X$  is the **moment of order 1**,

$$E(X) = \nu_1.$$

The **variance** of a random variable  $X$  is the **central moment of order 2**,

$$V(X) = \mu_2 = \nu_2 - \nu_1^2.$$

For any random variable  $X$ , the central moment of order 1 is 0,

$$\mu_1 = E(X - E(X)) = E(X) - E(X) = 0.$$

3. An important property of the moments of a random variable  $X$ , which we just state, without proof, is the following: If  $\underline{\nu}_n = \overline{E(|X|^n)}$  exists for some  $n \in \mathbb{N}$ , then  $\nu_k$ ,  $\underline{\nu}_k$  and  $\mu_k$  also exist, for all  $k = 1, n$ .